



**Institut de
radiophysique**
Rue du Grand-Pré 1
CH-1007 Lausanne

**Institut universitaire de
médecine sociale et préventive**
Route de la Corniche 2
CH-1066 Epalinges

Exposure of the Swiss population by Medical X-rays: 2008 Review

Abbas Aroua, Eleni T. Samara, François O. Bochud and Francis
R. Verdun

Institute of Radiation Physics, Lausanne

John-Paul Vader

University Institute of Social and Preventive Medicine, Lausanne

June 2011



This research project was jointly funded by the Swiss Federal Office of Public Health (contract Nr. 08.004670/434.0000/-53) and the Swiss National Science Foundation (subsidy Nr. 32003B_127461).

Abstract

Nationwide surveys on radiation dose to the population from medical radiology are recommended in order to follow the trends in population exposure and ensure radiation protection. The last survey in Switzerland was conducted in 1998 and the annual effective dose from medical radiology was then estimated to be 1 mSv/caput. The purpose of this work was to follow the trends in diagnostic and interventional radiology between 1998 and 2008 in Switzerland and determine the contribution of the different modalities and types of examinations to the collective effective dose from medical X-rays. An online database (www.raddose.ch) was developed to this effect, and all healthcare providers who hold a license to run an X-ray unit in the country were invited to participate in the survey. More than 225 examinations, covering eight radiological modalities, were considered and the average effective dose for each examination was reassessed. Data from about 3500 users were collected (42% response rate). The survey showed that the average frequency of X-ray examinations was 1.7 per caput and the average annual effective dose due to medical X-rays was 1.2 mSv/caput in 2008. The most frequent examinations were conventional and dental radiographies (88%). The contribution of computed tomography was only 6% in terms of examination frequency but as high as 68% in terms of the collective effective dose. The comparison with other countries showed that the collective effective dose in Switzerland was in the same range as in other countries with similar healthcare systems, although the annual number of examinations performed in Switzerland was higher.

Keywords

Diagnostic, interventional, radiology, medical radiation, X-rays, dose, population, survey

Zusammenfassung

Um die Entwicklung der öffentlichen Strahlenbelastung durch medizinische Radiologie zu verfolgen und einen entsprechenden Strahlenschutz gewährleisten zu können wird empfohlen landesweite Erhebungen zur Strahlendosis durch solche medizinischen Untersuchungen durchzuführen. Die letzte Erhebung in der Schweiz wurde 1998 durchgeführt, woraufhin die jährliche effektive Strahlendosis durch medizinische Radiologie auf 1 mSv pro Einwohner geschätzt wurde. Ziel dieser Arbeit ist Veränderungen der Strahlenbelastung durch diagnostische und interventionelle Radiologie zwischen 1998 und 2008 in der Schweiz zu untersuchen und den Beitrag von verschiedenen Methoden der medizinischen Röntgenuntersuchung zur gesamten effektiven Strahlendosis zu bestimmen. Für diesen Zweck wurde eine Online-Datenbank entwickelt (www.raddose.ch). Alle Gesundheitsversorger die über eine Lizenz für den Betrieb einer Röntgenanlage verfügen wurden gebeten, an der Erfassung teilzunehmen. Es wurden mehr als 225 Untersuchungen einbezogen, die 8 verschiedene radiologische Methoden abdecken. Hiermit wurde die mittlere effektive Dosis für eine Untersuchung neu bewertet. Insgesamt wurden Daten von 3500 Benutzern gesammelt, was einer Antwortquote von 42% entspricht. Die Erhebung zeigt, dass sich im Jahr 2008 im Mittel jeder Einwohner 1.7 Röntgenuntersuchung unterzogen hat und, dass die mittlere effektive Dosis in diesem Jahr 1.2 mSv pro Einwohner betrug. Die häufigsten Anwendungen (88%) waren konventionelle und zahnmedizinische Radiographie. Der Anteil der Computertomographie betrug bezüglich der Häufigkeit der Anwendung 6%, bezüglich der kollektiven effektiven Dosis aber 68%. Der Vergleich mit anderen Ländern zeigt, dass sich die kollektive effektive Strahlendosis in der Schweiz im selben Bereich wie die in Ländern mit ähnlichen Gesundheitssystemen befand. Die jährliche Zahl von Untersuchungen war in der Schweiz jedoch höher.

Résumé

Les enquêtes nationales visant à déterminer les doses de radiation délivrées à la population par la radiologie médicale sont recommandées afin de suivre les tendances de l'exposition de la population et assurer protection radiologique. La dernière enquête de ce genre a été menée en Suisse en 1998 et la dose efficace annuelle due à la radiologie médicale a été estimée à 1 mSv par habitant. Le but de ce travail était de suivre les tendances de la radiologie diagnostique et interventionnelle en Suisse, entre 1998 et 2008, et de déterminer la contribution des différentes modalités radiologiques et des divers types d'examens à la dose efficace collective due à l'application médicale des rayons X. Une base de données en ligne (www.raddose.ch) a été développée à cet effet, et tous les prestataires de soins détenant une autorisation à utiliser une installation radiologique dans le pays ont été invités à participer à l'enquête. Plus de 225 examens, couvrant huit modalités radiologiques, ont été considérés et la dose efficace moyenne pour chaque examen a été réévaluée. Les données concernant environ 3500 utilisateurs ont été recueillies, correspondant à un taux de réponse de 42%. L'enquête a montré qu'en 2008, la fréquence moyenne des examens radiodiagnostiques était de 1,7 par habitant et la dose efficace annuelle moyenne due aux rayons X médicaux était de 1,2 mSv par habitant. Les examens les plus fréquents étaient les radiographies conventionnelles et dentaires (88%). La contribution de la tomодensitométrie a été de 6% en termes de fréquence d'examens, et de 68% en termes de dose efficace collective. La comparaison avec d'autres pays a montré que la dose efficace moyenne par habitant en Suisse était dans la même gamme que celle enregistrée dans d'autres pays avec des systèmes de santé publique similaires, bien que le nombre annuel d'examens effectués en Suisse a été plus élevé à cause de la contribution élevée de la radiologie dentaire.

Contents

Introduction6
Methods.....7
Results and discussion..... 11
Conclusion..... 28
References..... 29
Appendices 33

Introduction

Medical exposure to X-rays represents a major source of man-made irradiation of the population. At the global level, the total annual per caput effective dose, due to all sources of irradiation, is about 3.1 mSv [UNSCEAR 2010]. Although diagnostic radiology represents only 20% of it, it accounts for more than 94% of the man-made component. Similarly, in Switzerland where the annual per caput effective dose amounted to 4 mSv in 2009 [FOPH 2010], diagnostic radiology represented 30% of the total, but more than 92% of man-made irradiation.

This is why both at national and international levels, surveying the population exposure by medical X-rays is recommended as a useful tool in radiation protection. While the Swiss Federal Act on Radiation Protection stipulates in Art. 5 that “the Confederation shall encourage scientific research on the effects of radiation and on radiation protection” [LRaP 2004], the European Commission states that the “Member States shall ensure that the distribution of individual dose estimates from medical exposure is determined for the population and for relevant reference groups of the population as may be deemed necessary by the Member State” [EU 1997]. Among the main objectives of population dose assessments are: “1) to observe trends in the annual collective dose and the annual average per caput dose from medical x-rays in a country with time; 2) to determine the contributions of different imaging modalities and types of examination to the total collective dose from all medical x-rays.” [EC 2008a]

Because of the pace of technological developments in the field of radiology and the evolution of the medical practice, the situation is reassessed ideally every five years as recommended by the European Guidance on Estimating Population Doses from Medical X-Ray Procedures [EC 2008a, EC 2008b], or at least every ten years. At the international level, surveys on the exposure of the global population by medical radiology are conducted every decade by the United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR). UNSCEAR has just issued its 2008 report [UNSCEAR 2010] with data covering the period 1997-2007.

Recently, several countries conducted nationwide surveys including Finland [Järvinen 2011] (2008 data), France [IRSN-IVS 2010] (2007 data), Germany [Bernhard-Ströl et al. 2010] (2008 data), the Netherlands [RIVM 2010] (2008 data), Norway [NRPA 2010] (2008 data), the United Kingdom [HPA 2010] (2008 data), and the United States of America [NCRP 2009] (2006 data).

Switzerland started surveying the exposure of the population by medical X-rays back in the late 1950s [Zuppinger et al. 1961, Poretti et al. 1971, Mini and Poretti 1984, Mini 1992]. The exposure of the Swiss population is reviewed periodically, every decade

with a full re-evaluation survey addressing all healthcare providers and every five years with an updating survey covering a small stratified sample of healthcare providers. The last full re-evaluation survey concerned the 1998 data [Aroua et al. 2002a, Aroua et al. 2002b]. It provided a significant amount of information on the frequency of the X-ray examinations performed in Switzerland and the associated radiation doses. The last updating survey concerned the 2003 data [Aroua et al. 2007a]. It revealed, for example, that the use of computed tomography (CT) had registered a sharp increase (70%) in a 5-year period.

The aim of the present work was to reassess the exposure of the Swiss population to medical X-rays in 2008 and to update the associated annual frequency of examinations, the effective doses per examination and the annual collective effective dose.

Methods

The survey was led by a steering group comprising representatives of Federal Office of Public Health (FOPH), the Institute of Radiation Physics (IRA) from the University Department of Radiology (DRM – CHUV, Lausanne) and the University Institute of Social and Preventive Medicine (IUMSP) from the University Department of Community Medicine and Health (DUMSC – CHUV, Lausanne). It was followed by a support group comprising representatives of most scientific and professional societies concerned: Association Romande de Radioprotection, College of General Practitioners, Federal Commission for Radiation Protection and Radioactivity Surveillance, Santésuisse, Swiss Society of Cantonal Chief-Physicians, Swiss Society of Cardiology, Swiss Society of Chiropractic, Swiss Society of Dental Medicine, Swiss Society of Gastroenterology, Swiss Society of General Medicine, Swiss Society of Internal Medicine, Swiss Society of Paediatrics, Swiss Society of Radiobiology and Medical Physics, Swiss Society of Radiographers, Swiss Society of Radiology, Swiss Society of Urology, and The Swiss Hospitals.

The recommendations of the European Group “Dose Datamed” on how to conduct national surveys on population exposure, which were accepted by the European Commission [EC 2008a], were followed in this survey.

During the 1998 nationwide survey all Swiss healthcare providers were addressed except general and dental medicine practices which were sampled at rates of 20% and 10% respectively. This time the whole community of healthcare providers possessing an authorisation to run an X-ray unit was addressed. This amounts to 8247 practices, radiology institutes and hospital departments (Table 1), running 17391 X-ray units of all kinds (Table 2) authorised by the Regulatory Authority which, in Switzerland, is the Federal Office of Public Health. The 8247 healthcare providers are distributed into 4587

medical, 3526 dental and 134 chiropractic. They cover 7779 practices (individual and group), 383 hospital departments and 85 radiology institutes.

Table 1: Healthcare providers performing X-rays in Switzerland

Category of healthcare provider	Number
Individual medical practices	3568
Group medical practices	529
University hospitals	53
Cantonal hospitals	62
District hospitals	111
City hospitals	14
Private hospitals	132
Homes for the elderly	11
Radiology institutes	85
Others	22
Total medical	4587
Individual chiropractic practices	125
Group chiropractic practices	9
Total chiropractic	134
Individual dental practices	3072
Group dental practices	284
Dental clinics	13
School dental radiology centres	124
Others	33
Total dental	3526
Grand total	8247

Table 2: Various types of X-rays units used in Switzerland

	Use of X-ray unit	Number
Total		17391
Generic	Dental	10553
	Medical	6704
	Chiropractic	134
Specific	Radiography	4937
	Radiography + Fluoroscopy	760
	Fluoroscopy (mobile)	476
	Interventional (intensive dose)	127
	Computed tomography	238
	Mammography	249
	Bone densitometry	147
	Small dental (< 70 kV)	8572
	Orthopantomography	1171
	Panoramic + teleradiography	661
	Teleradiography	4
Dental volume tomography	49	

In order to increase their probability of participation, the participants were offered several options on how to provide their frequency data: paper form by post mail, electronic form by email or by registration online. For this purpose, a dedicated website (raddose.ch) was developed beforehand (see Appendix 1). In the first letter sent to them, the participants received their own username and password to access their form.

The participants were encouraged to fill in a form with reference categories of examinations, based on the “Dose Datamed” methodology [EC 2008a]. This consists in 225 examination types grouped in seven radiological modalities: radiography, conventional fluoroscopy, diagnostic interventional radiology, therapeutic interventional radiology, computed tomography, dental radiology, mammography, and bone densitometry. In the case the participant was not able to provide data concerning the 225 examination types, he/she could use 70 broader categories (see Appendix 2). The participants were also allowed to send their data in the format of their choice: local categories or medical tariff codes (Tarmed).

After ten months and three reminders the frequency data collection was closed and the data processing started: introducing in the database the statistics received by email or in paper form, checking the data registered online and correcting the typing errors where necessary. The data received in local categories or in Tarmed format were redistributed over the reference categories.

National frequencies were established by projecting the collected statistics according to the number of X-ray units in the survey, with the following formula:

$$N_{2008} = \sum_{i,j} \frac{I_{total}(i,j)}{I_{part}(i,j)} \cdot N_{part}(i,j)$$

where N_{2008} is the national examination frequency, i refers to the healthcare provider (medical practice, dental practice, chiropractor, hospital department, radiology institute), j refers to the type of X-ray unit, I_{total} being the number of X-ray units in Switzerland, I_{part} the number of X-ray units run by healthcare providers who participated in the survey, and N_{part} the number of the examinations performed by the healthcare providers who participated in the survey.

Unlike the 1998 survey where the number of practices and hospital departments was used to project the data associated with the participating sample to the total number in the country, in the present investigation the number of X-ray units was available and we chose to use it since it leads to more accurate results. The difference in the frequency of examinations obtained by the two approaches was explored.

In the 1998 survey the effective doses related to the various types of examinations were calculated using dosimetric models specific to each radiological modality [Aroua et al. 2002b]. The effective doses were updated the last couple of years by surveying some fluoroscopic procedures in large and small hospitals [Aroua et al. 2007b, Samara et al. 2010, Samara et al. 2011]. For radiography examinations, the technical parameters were reassessed and the effective doses were calculated using the software program PCXMC [Tapiovaara et al. 1997] (See Appendix 3). The 2008 version of PCXMC was adopted (12, PCXMC version 2.0, 2008, STUK, Helsinki, Finland). For CT examinations, the effective doses were reviewed in an auditing campaign [Treier et al. 2010] (See Appendix 3). For some other examinations, bibliographic data were used [Brambilla et al. 2004, Chamberlain et al. 2000, Crawley and Rogers 2000, Crawley 2004, Danforth and Clark 2000, Delichas et al. 2004, Gibbs 2000, Gijbels et al. 2004, Gijbels et al. 2005, Hatzioannou et al. 2000, Karthikesalingam et al. 2009, Ludlow et al. 2008, Martin 2008, Mettler et al. 2008, Pantos et al. 2009, Perisinakis et al. 2003, Perisinakis et al. 2004, Ruiz Cruces et al. 1998, Smith et al. 2009, Şorop et al. 2008, Thomas et al. 2006, Tsalafoutas et al. 2005, Yakoumakis et al. 2001].

In the calculation of the updated effective doses the tissue weighting factors (w_T) given by the International Commission on Radiological Protection in its Publication 60 [ICRP 1991] were used. A couple of years ago the ICRP published new values in Publication 103 [ICRP 2007] as shown in Table 3.

The main changes are a decrease of the w_T of the gonads by a factor 0.4 and an increase of that of the breast by a factor 2.4. The two sets of w_T were compared for two radiological modalities. In the case of radiography the new figures lead to a 14% decrease in the collective dose, whereas in the case of CT a 2% increase is registered (see Appendix 4). In the most recent investigation performed in the UK [HPA 2010] a 2-3% decrease in the collective dose was registered for conventional radiology including dental and for CT and a 12-13% increase for interventional radiology. This was reflected in an overall 2% increase of the total collective dose due the w_T change.

Table 3: Tissue weighting factors (w_T)

Organ/Tissue	ICRP ₆₀	ICRP ₁₀₃	ICRP ₁₀₃ /ICRP ₆₀
Gonads	0.20	0.08	0.4
Colon	0.12	0.12	1.0
Lungs	0.12	0.12	1.0
Red bone marrow	0.12	0.12	1.0
Stomach	0.12	0.12	1.0
Bladder	0.05	0.04	0.8
Breast	0.05	0.12	2.4
Liver	0.05	0.04	0.8
Oesophagus	0.05	0.04	0.8
Thyroid	0.05	0.04	0.8
Bone surface	0.01	0.01	1.0
Skin	0.01	0.01	1.0
Brain		0.01	
Salivary glands		0.01	
Remainder	0.05*	0.01 ⁺	

* ICRP60 remainder tissues/organs: adrenals, brain, kidneys, muscle, pancreas, small intestine, large intestine, spleen, thymus, uterus.

⁺ ICRP103 remainder tissues/organs: adrenals, extrathoracic tissue, gall bladder, heart, kidneys, lymphatic nodes, muscle, oral mucosa, pancreas, prostate, small intestine, spleen, thymus, uterus/cervix.

Results and discussion

The rate of return (address change, retired, deceased) was about 1% and the rate of explicit refusal was about 0.2%. Table 4 shows the response rates for the three broad categories of health care providers (medical, dental and chiropractic) as well as for various types of hospital departments.

In terms of number of healthcare providers, the overall response rate is 42% corresponding to 3486 respondents. In terms of X-ray units, the overall response rate is 45%, corresponding to 7878 units.

During the 1998 survey, a sample of 3000 establishments was contacted and at the end of the survey the response rate was 60% [Aroua et al. 2002a]. About 1800 participants sent their data, which is nearly half the number of respondents in the 2008 survey.

Two participants over three registered their data online. Half of the rest sent their data in electronic form and the other half in paper form.

Table 4: Response rates for the 2008 Swiss nationwide survey

Category of healthcare providers	Healthcare providers			X-ray units		
	Total	Respondents	Rate (%)	Total	Respondents	Rate (%)
All categories	8247	3486	42	17391	7878	45
Medical	4587	1953	43	6704	3376	50
Dental	3526	1461	41	10553	4418	42
Chiropractic	134	89	66	134	89	66
Radiology institutes	85	39	46	347	170	49
Hospital departments (all)	383	249	65	2394	1519	63
in university hospitals	53	26	49	543	242	45
in canton hospitals	62	40	65	481	354	74
in district hospitals	111	80	72	685	446	65
in state hospitals	14	10	71	94	67	71
in private hospitals	132	87	66	576	403	70
in houses for the elderly	11	6	55	15	7	47

Table 5 gives the range of regional variation of the response rates. The seven regions of Switzerland are presented in Figure 1. The cantons that constitute each region are given in table 6. If all categories are summed up, the regional variation is in the range 41-46%. The detailed regional response rates are given in Appendix 5.

Table 5: Regional distribution of the response rates (%) for the 2008 Swiss nationwide survey. CH : Whole Switzerland.

Category of health care providers	Total #	Region							CH
		1	2	3	4	5	6	7	
All categories	8247	40	43	44	41	45	42	46	42
Medical	4587	40	43	43	42	45	41	45	43
Dental	3526	40	42	44	37	43	42	47	41
Chiropractic	134	50	66	64	81	87	63	100	66
Radiology institutes	85	35	53	50	73	27	100	33	46
Hospital departments (all)	383	60	63	71	64	69	59	77	65
in university hospitals	53	43	36	57	78	—	—	—	49
in canton hospitals	62	—	60	83	50	60	73	—	65
in district hospitals	111	59	79	78	67	74	50	83	72
in state hospitals	14	—	40	100	86	—	—	—	71
in private hospitals	132	74	55	62	43	80	43	83	66
in houses for the elderly	11	—	60	—	100	—	—	50	55



Figure 1: The Seven region of Switzerland according to the Swiss Federal Office of Statistics

Table 6: The Cantons included in the seven regions of Switzerland

Region	Cantons
1. Lake Geneva	VD / VS / GE
2. Espace Mittelland	BE / FR / SO / NE / JU
3. Northwestern Switzerland	BS / BL / AG
4. Zurich	ZH
5. Eastern Switzerland	GL / SH / AR / AI / SG / GR / TG
6. Central Switzerland	LU / UR / SZ / OW / NW / ZG
7. Ticino	TI

A sensitivity analysis was performed in order to assess the effect on the result if the non respondents had features that are different from those of the respondents, in particular the frequencies of examinations performed. The hypothesis that the non-participants perform 20% more or 20% less examinations than the participants resulted in an average 10% difference in the national examination frequencies. As shown in Appendix 6, the difference is smaller for interventional radiology (-3%, +4%) and bigger for dental radiology (-12%, +12%). For dental radiology, the number of non-participants is higher than that of participants.

Table 7 shows the annual number of examinations performed in Switzerland in 2008 (7.7M population), the number per thousand population, as well as the associated annual collective dose and the average per caput effective dose delivered by the whole medical radiodiagnostics as well as by the various radiological modalities. The detailed frequency and dose results are provided in Appendix 9.

Table 7: 2008 Swiss annual frequency and dose data

Radiological modality	Number of examinations (in thousands)	Collective dose (man.Sv)	Number of examinations per 1000 population	Effective dose per caput (mSv)
Radiography	6000	1330	780	0.17
Conventional fluoroscopy	153	415	20	0.05
Interventional – diagnostic	56	553	7.2	0.07
Interventional – therapeutic	46	528	6.0	0.07
Computed tomography	780	6150	100	0.8
Dental radiology	5430	63	700	0.01
Mammography	387	62	50	0.01
Bone densitometry	117	0.31	15	0.00004
Total	13'000	9100	1700	1.2

This investigation revealed that the total number of X-ray examinations carried out in Switzerland was 13 M per year in 2008, corresponding to 1.7 examinations per caput. The associated collective effective dose accounts to about 9100 man Sv which, reported to the Swiss population gives an average effective dose of 1.2 mSv per caput. In the 2003 review the average effective dose was also estimated to 1.2 mSv per caput per year. Therefore it registered no change although the frequency increased during this 5-year period. This is due to the change of the dose vector. In fact the updated effective doses per examination for radiographies are significantly lower than older values. The use of the dose vector established in 1998 with the 2008 frequency data would have resulted in an average effective dose of 1.4 mSv per caput.

Figure 2 presents the distribution of the total annual number of examinations and the total annual collective dose over the various radiological modalities. It shows that the highest contributions to the total number of examinations come from radiography (46%) and dental radiology (42%). However, in terms of the collective effective dose the contribution of radiography drops to 14.6% and that of dental radiology to less than 1%. On the contrary CT which contributes 6% to the number of examinations is responsible of more than 2/3 of the collective effective dose. Similarly, interventional radiology (diagnostic and therapeutic) whose contribution in terms of frequency of examinations is only 0.8% delivers nearly 12% of the collective effective dose.





Figure 2: Distribution of the total annual number of examinations (upper part) and the total annual collective dose (lower part) over the various radiological modalities: radiography (RA), conventional fluoroscopy (FL), diagnostic interventional radiology (ID), therapeutic interventional radiology (IT), computed tomography (CT), dental radiology (DR), mammography (MA), bone densitometry (BD).

Figures 3 to 6 present the 3 most frequent examinations and the 3 examinations with the highest dose contribution in radiography, interventional radiology, computed tomography, and dental radiology.

RA / Frequency



RA / Dose





Figure 3: The 3 most frequent examinations and the 3 examinations with the highest dose contribution in radiography

IN / Frequency



IN / Dose





Figure 4: The 3 most frequent examinations and the 3 examinations with the highest dose contribution in interventional radiology (both diagnostic and therapeutic)

CT / Frequency



CT / Dose





Figure 5: The 3 most frequent examinations and the 3 examinations with the highest dose contribution in computed tomography

DR / Frequency



DR / Dose




Figure 6: The 3 most frequent examinations and the 3 examinations with the highest dose contribution in dental radiology

Figure 7 shows the frequency and dose contribution of the different healthcare providers (The details are given in Appendices 7 and 8). Hospitals undertake 34.6% of the examinations and are responsible of 81% of the collective effective dose. Radiology institutes contribute 4% in terms of frequency and 12.6% in terms of the collective effective dose. Dental practices perform almost 41% of the examinations and are responsible of only 0.7% of the collective dose. Medical practices perform a fifth of the examinations and contribute 5.3% to the collective effective dose. Chiropractors are responsible for 0.5% of the frequencies and 0.5% of the collective effective dose.




Figure 7: Frequency (upper part) and dose (lower part) contribution of the different healthcare providers

Tables 8 and 9 present the frequency and dose contribution of the different healthcare providers for the 8 radiological modalities considered.

Table 8: Frequency contribution of the different healthcare providers (%) for the 8 radiological modalities considered

	Dental practices	Medical practices	Chiropractors	Hospital departments	Radiology institutes
RA	0	42.5	1.1	52.8	3.7
FL	0	6.1	0	77.1	16.8
ID	0	0	0	99.9	0.1
IT	0	0	0	97.5	2.5
CT	1.5	0	0	83.3	15.2
DR	97.2	0.2	0	2.4	0.1
MA	0	7.7	0	61.5	30.8
BD	0	2.4	0	75.9	21.7

Table 9: Dose contribution of the different healthcare providers (%) for the 8 radiological modalities considered

	Dental practices	Medical practices	Chiropractors	Hospital departments	Radiology institutes
RA	0	34	3.7	55	6.4
FL	0	5.1	0	87	8.0
ID	0	0	0	100	0.05
IT	0	0	0	100	0.003
CT	0.1	0	0	83.5	16.4
DR	92.0	2.1	0	5.1	0.8
MA	0	7.7	0	61.5	30.8
BD	0	2.4	0	77.5	20.2

Table 10 compares the frequency and dose data obtained in the present study with the data established in Switzerland in 1998.

Table 10: 2008/1998 ratios of the frequency and dose in Switzerland

Radiological modality	Number of examinations	Collective dose	Number of examinations / 1000 population	Effective dose / caput
Radiography	1.32	0.45	1.21	0.42
Conventional fluoroscopy	0.98	0.34	0.90	0.31
Interventional – diagnostic	0.79	1.00	0.73	0.93
Interventional – therapeutic	1.72	2.06	1.59	1.90
Computed tomography	2.38	3.11	2.19	2.86
Dental radiology	1.32	0.87	1.21	0.80
Mammography	1.73	1.43	1.60	1.32
Bone densitometry	3.64	7.55	3.36	6.96
Total	1.36	1.30	1.26	1.20

The first two columns of the table present the ratio of examination frequency and collective effective dose relative to 2008 and 1998 (absolute values) and the last two columns the ratio of the number of examinations per 1000 population and the average effective dose per caput. Some of the increase in the absolute values may be associated with the increase in the Swiss population (8.5% in a decade). Thus, to eliminate the demographic factor, the examination frequencies per 1000 population and the average effective doses per caput were compared.

Unlike the 1998 survey where the number of practices and hospital departments was used to project the data associated with the participating sample to the total number in the country, in the present investigation the number of X-ray units was available and we chose to use it since it leads to more accurate results. This different approach may also explain some of the differences between the examination frequencies in 1998 and 2008 surveys, i.e. the 21% increase in conventional and dental radiographies.

The number of radiographies registered a 20% increase but the average effective dose per caput decreased by more than a factor 2. This is due the significant reduction in the average effective dose per radiography for many types of examinations.

The number of conventional fluoroscopy and diagnostic interventional procedures showed a decrease of 10% and 30%, respectively, in terms of number of examinations. Inversely, the number of therapeutic intervention procedures showed an increase of 60% in a decade, corresponding to an increase of effective dose per caput of 90%. This

can be explained by the fact that cases previously treated with open surgery may be now safely and effectively treated by interventional procedures [Balter et al. 2008]. Accordingly, the increase in the dose delivered by this kind of procedures may be attributed to the fact that more complex cases may now be treated by interventional radiology, resulting, however, in higher patient doses [Balter et al. 2008].

The main increase in the collective effective dose in a decade may be attributed to the increase in CT (286% in terms of the collective effective dose since 1998). The increase in the number of CT examinations had already been observed [Aroua et al. 2007a] and attributed to the technology advance in CT scanners that led to the change of medical practice by replacing fluoroscopy guided procedures with CT scans. This may also explain the reduction in the number of diagnostic interventional procedures. It is important to note here that for this survey no CT scans associated to SPECT/CT examinations, PET/CT examinations or for radiation therapy planning were taken into account, as they will be considered in a specific survey dedicated to the dose delivered in nuclear medicine [Roser 2011].

The number of bone densitometry examinations has notably increased (3.36 times) since 1998; however, its associated dose was too low to significantly affect the collective effective dose (see Table 7). Some increase in these examinations may be attributed to the population ageing but also to the fact that bone densitometry examinations are performed for the diagnosis of osteoporosis as well as the follow-up of osteoporosis treatments.

Table 11 presents the number of all examinations and number of CT examinations (per 1000 population), average effective dose due to radiodiagnostics and average effective dose due to CT, for 1998, 2003 and 2008. Both the frequency of CT examinations and the associated collective effective dose registered a steady increase since 1998: respectively a factor of 2.2 and 2.9 in a decade. It should be noted that the increase was higher between 1998 and 2003 than between 2003 and 2008. In one decade, the contribution of CT to the total medical X-rays increased from 3.4% to 6% in terms of the frequency and from 28% to 68% in terms of the collective effective dose.

Table 11: Number of all examinations and number of CT examinations (per 1000 population), average effective dose due to radiodiagnostics and average effective dose due to CT, for 1998, 2003 and 2008 (absolute and relative values)

Year	N/1000 population		CT /Total (%)	E (mSv/caput)		CT /Total (%)
	Total	CT		Total	CT	
1998	1340	46.2	3.4	1.0	0.28	28
2003	1470	76.7	5.2	1.2	0.56	47
2008	1680	100	6.0	1.2	0.80	68

Ratio	N/1000 population		E (mSv/caput)	
	Total	CT	Total	CT
2003/1998	1.10	1.66	1.20	2.00
2008/2003	1.14	1.32	1.00	1.43
2008/1998	1.26	2.19	1.20	2.86

Table 12 shows the evolution of the number of X-ray Units in Switzerland from 1998 and 2008. The increase in the number of CT scanners, and dental and mammography X-ray units in a decade (27%, 22% and 4% respectively) may explain the part of the increase in the frequency of examinations for those radiological modalities.

Table 12: Number of X-ray Units in Switzerland

Specialty	Radiological modality	Code	1998*	2008**	08/98
Medical	Radiography	RA	—	4937	
	Fluoroscopy (mobile)	RD	—	476	
	Radiography & fluoroscopy	RAD	—	760	
	Radiography & fluoroscopy (Intensive Dose)	RDI	—	127	
	Computed tomography	CT	187	238	1.27
	Mammography	MAM	240	249	1.04
	Bone densitometry	KDM	—	147	
Total "Medical"		—	8419	6934	0.82
Dental	Intra-oral (< 70 kV)	RKL	—	8572	
	Orthopantomography	OPG	—	1171	
	Teleradiography	FR		4	
	Orthopantomography & Teleradiography	OPF	—	661	
	Dental volume tomography	DVT	—	49	
Total "Dental"		—	8583	10457	1.22
Total "Medial + Dental"		—	17002	17391	1.02

* UNSCEAR 2000

** BEBERA file received from FOPH

Table 13 shows the contribution of medical exposure to the total irradiation of the population for various countries. It ranges from about 15% in Finland to as high as 48% in the USA.

Table 13: Contribution of medical exposure to the total irradiation of the population

Country	CH	NO	D	USA	UK	FI
E (mSv/caput)	4.0	4.4	3.9	6.2	2.7	3.7
Contributions (%)						
Radon	40	46	28	37	49	54.2
Medicine	30	24	46	48	16	14.4
	NM	OUT	IN	IN	IN	OUT
Cosmic	10	7	7.7	5	12	8.9
Terrestrial	9	14	10	3	13	12.7
Internal	9	7	7.7	5	9	9.8
Others	2	1	0.5	2	1	—

Table 14 compares the frequency and dose data obtained in the present study with the data reported recently by seven countries: Finland, France, Germany, the Netherlands, Norway, the UK and the USA. The data presented in the 2008 UNSCEAR report for countries with similar healthcare systems as Switzerland (level I), which covers the decade 1997-2007, is also included for comparison.

Table 14: Frequency and dose comparisons with data reported in other countries

Country	Number of X-ray examinations per 1000 population	Annual per caput effective dose due to medical X-rays (mSv)
UNSCEAR – Health level I (1997-2007)	1607	1.9
USA (2006)	1257	2.2
France (2007) *	1152	1.2
Germany (2008)	1650	1.7
United Kingdom (2008)	752	0.4
The Netherlands (2008)	573	0.7
Norway (2008) **	670	1.1
Finland (2008)	717	0.45
Switzerland (2008)	1700	1.2

* Excluding therapeutic interventional procedures – ** Excluding dental radiology

For the Netherlands, France and the USA nuclear medicine was included in the originally reported figures and had to be removed in Table 14 for the comparison of the X-ray component.

The annual number of examinations per 1000 population established in this work (1700) is the highest among the countries of interest. This is due to the high frequency of dental X-ray examinations in Switzerland (see also Table 18). The average annual effective dose per caput of 1.2 mSv compares well with the figures reported in other countries ranging from 0.4 mSv in the UK to 2.2 mSv in the USA.

Table 15 gives nuclear medicine frequency and dose comparisons with data reported in other countries. The Swiss frequency and dose values are relatively low and compare with those registered in Norway. They are well below the figures reported in the USA. If we consider the contribution of nuclear medicine to the total average effective dose (radiodiagnostics + nuclear medicine), one can observe the existence of three categories of countries with NM contributions amounting to 5% such as Switzerland, Germany and the Scandinavian countries, 10% such as France and the Netherlands, and 30% in the case of the USA.

Table 15: Nuclear medicine (NM) frequency and dose, and total dose, comparisons with data reported in other countries

Country	Number of NM procedures per 1000 population	Annual per caput E due to NM (mSv)	Total annual per caput E (mSv)	E ratio NM / Total (%)
UNSCEAR – Health level I (1997-2007)	22.1	0.12	2.0	6
USA (2006)	60	0.8	3.0	27
France (2007)	18	0.13	1.3	10
Germany (2008)	37	0.1	1.8	5.6
The Netherlands (2008)	24	0.095	0.8	12
Norway (2008) *	9.7	0.054	1.1	4.9
Finland (2008)	7.7	0.03	0.48	6.3
Switzerland (2010) **	13	0.06	1.3	4.6

* [Frede Unhjem 2011] – ** [Roser 2011]

Table 16 compares the contribution of CT to the total number of examinations and to the collective effective dose in Switzerland with that reported in other countries. It shows clearly that the same pattern observed in Switzerland is registered elsewhere: a 10-20% contribution in terms of frequencies is reflected into up to a 2/3 contribution in terms of collective effective dose. In the case of Norway the CT frequency contribution is even higher (29%), since dental radiology is not considered.

Table 16: Contribution of computed tomography (%)

Country	To the total number of X-ray examinations	To the total collective dose
UNSCEAR – Health level I (1997-2007)	8	47
USA (2006)	18	66
France (2007) *	10	65
Germany (2008)	8	60
United Kingdom (2008)	7	67
The Netherlands (2008)	11	53
Norway (2008) **	29	80
Finland (2008)	8.3	58
Switzerland (2008)	6	68

* Excluding therapeutic interventional procedures – ** Excluding dental radiology.

Table 17 compares the contribution of interventional radiology to the total number of examinations and to the collective effective dose in Switzerland with that reported in other countries. The Swiss frequency contribution data (about 1%) compare with those of other countries except the USA and the UK where the figures are relatively high (5% and 12% respectively) and Norway where the figure is relatively low (0.2%). The contribution of interventional radiology to the total collective effective dose ranges from 6% in France (where therapeutic procedures are excluded) to nearly 20% in Germany and the USA. The Swiss figure (12%) compares with that other countries.

Table 17: Contribution of interventional radiology both diagnostic and therapeutic (%)

Country	To the total number of X-ray examinations	To the total collective effective dose
UNSCEAR – Health level I (1997-2007)	0.5	4.3
USA (2006)	5	19
France (2007) *	0.6	6.1
Germany (2008)	2	19
United Kingdom (2008)	12	13
Norway (2008) **	0.2	10
Finland (2008)	0.8	14
Switzerland (2008)	0.8	12

* Excluding therapeutic interventional procedures – ** Excluding dental radiology [Almén 2011]

Table 18 compares the frequency and dose data obtained in the present study for dental radiology with the data reported recently by seven countries. The total annual number of examinations per 1000 population established in this work (1700) is higher than the figures of the other countries. This is due to the high frequency of dental X-ray examinations in Switzerland: 700 per year per 1000 population, compared to 289 in France and 193 in the UK. The US frequency figure for dental radiology seems to be too low.

Table 18: Dental radiology frequency and dose comparisons with data reported in other countries (%)

Country	Number of dental examinations per 1000 population	Annual per caput E due to dental radiology (microSv)	% of total number of X-ray examinations	% of total collective dose
UNSCEAR – Health level I (1997-2007)	275	6	17	0.3
USA (2006)	8.8	8	0.7	0.4
France (2007)	289	3	25	0.3
Germany (2008)	600	≤10	37	0.3
United Kingdom (2008)	193	2	26	0.5
Norway (2008) *	1173	26.2	64	2.3
Finland (2008)	385	3	30	0.7
Switzerland (2008)	700	8	42	0.7

* [NRPA 2009]

Conclusion

This work updated the frequency and collective effective dose data for medical and dental X-ray examinations in Switzerland for the year 2008. The average frequency of X-ray examinations was found to be 1.7 per caput with an associated average annual effective dose of 1.2 mSv/caput. There was no increase of the average effective dose since 2003 although the frequency has increased due to the fact that the effective doses per examination decreased significantly for radiography. The survey showed that Switzerland stands at the same level as other countries with similar healthcare systems in terms of collective effective dose. However, the annual number of examinations per capita in Switzerland was higher than that performed in other countries, due to the high number of dental examinations performed in the country. The main contributor to the collective effective dose was computed tomography. It is important to regularly perform surveys in order to follow the trends in population exposure. In addition, such surveys, along with the definition and implementation of diagnostic reference levels may identify the points to focus for future optimisation campaigns.

References

- [Almén 2011] Almén A. Personal communication; 2011
- [Aroua et al. 2002a] Aroua A, Burnand B, Decka I, Vader JP and Valley JF. Nationwide Survey on Radiation Doses in Diagnostic and In-terventional Radiology in Switzerland in 1998. *Health Phys* 83(1):46-55; 2002
- [Aroua et al. 2002b] Aroua A, Decka I, Burnand B, Vader JP and Valley JF. Dosimetric aspects of a national survey of diagnostic and interventional radiology in switzerland. *Medical Phys* 29(10):2247:2259; 2002
- [Aroua et al. 2007a] Aroua A, Vader JP, Valley JF and Verdun FR. Exposure of the Swiss population by radiodiagnostics: 2003 review. *Health Phys* 92(5):442-8; 2007
- [Aroua et al. 2007b] Aroua A, Rickli H, Stauffer JC, Schnyder P, Trueb Ph, Valley JF, Vock P, Verdun FR. How to set up and apply reference levels in fluoroscopy at a national level? *European Radiology* 17(6):1621-33; 2007
- [Balter et al. 2008] For the ref. Balter S, Miller DL, Vano E, Ortiz Lopez P, Bernardi G, Coteló E, Faulkner K, Nowotny R, Padovani R, Ramirez A. A pilot study exploring the possibility of establishing guidance levels in x-ray directed interventional procedures. *Med Phys*. 35:673-80; 2008.
- [Bernhard-Ströl et al. 2010] Bernhard-Ströl C, Hachenberger C, Trugenberger-Schnabel A, Peter J. Jahresbericht 2009 Umweltradioaktivität und Strahlenbelastung. Bundesamt für Strahlenschutz Salzgitter; 2010
- [Brambilla et al. 2004] Brambilla M, Marano G, Dominietto M, Raffaele A, Carriero A. Patient radiation doses and references levels in interventional radiology. *La Radiologia Medica*. 408-418; 2004
- [Chamberlain et al. 2000] Chamberlain CC, Huda W, Hojnowski LS, Perkins A, Scaramuzzino A. Radiation doses to patients undergoing scoliosis radiography. *The British journal of radiology*. 73(872):847-853; 2000
- [Crawley and Rogers 2000] Crawley MT, Rogers AT. Dose-area product measurements in a range of common orthopaedic procedures and their possible use in establishing local diagnostic reference levels. *The British journal of radiology*. 73(871):740-744; 2000
- [Crawley 2004] Crawley MT. Patient and operator dose during fluoroscopic examination of swallow mechanism. *British Journal of Radiology*. 77(920):654-656; 2004
- [Danforth and Clark 2000] Danforth R, Clark D. Effective dose from radiation absorbed during a panoramic examination with a new generation machine Oral surgery, oral medicine, oral pathology, oral radiology and endodontics 103(3):437-440; 2000
- [Deak et al. 2010] Deak PD, Smal Y, Kalender WA. Multisection CT protocols: sex- and age-specific conversion factors used to determine effective dose from dose-length product. *Radiology*. 257:158-66; 2010.
- [Delichas et al. 2004] Delichas MG, Hatziiannou K, Papanastassiou E, Albanopoulou P, Chatzi E, Sioundas A, Psarrakos K. Radiation doses to patients undergoing barium meal and barium enema examinations. *Radiation protection dosimetry*. 109(3):243-247; 2004
- [EC 2008a] European Commission. European Guidance on Estimating Population Doses from Medical X-ray Procedures. Radiation protection no. 154. Brussels: Directorate General for Energy and Transport; 2008
- [EC 2008b] European Commission. Review of Recent National Surveys of Population Exposure from Medical X-rays in Europe. Annex to the European Guidance on

- Estimating Population Doses from Medical X-ray Procedures. Radiation protection no. 154. Brussels: Directorate General for Energy and Transport; 2008
- [EU 1997] Council Directive 97/43/Euratom of 30 June 1997. The Medical Exposures Directive Official Journal 180:22-27; 1997
- [FOPH 2010] Federal Office of Public Health. 2009 Annual Report of the Radiological Protection Division. Bern; 2010
- [Gibbs 2000] Gibbs SJ. Effective dose equivalent and effective dose: comparison for common projections in oral and maxillofacial radiology. Oral surgery, oral medicine, oral pathology, oral radiology and endodontics. 90(4):538-45; 2000
- [Gijbels et al. 2004] Gijbels F, Sanderink G, Wyatt J, Van Dam J, Nowak B, Jacobs R. Radiation doses of indirect and direct digital cephalometric radiography. British dental journal. 197(3):149-52; 2004
- [Gijbels et al. 2005] Gijbels F, Jacobs R, Bogaerts R, Debaveye D, Verlinden S, Sanderink G. Dosimetry of digital panoramic imaging. Part I: Patient exposure. Dento maxillo facial radiology. 34(3):145-91; 2005
- [Hatzioannou et al. 2000] Hatzioannou KA, Psarrakos K, Molyvda-Athanasopoulou E, Kitis G, Papanastassiou E, Sofroniadis I, Kimoundri O. Dosimetric considerations in mammography. Eur Radiol. 10:1193-1196; 2000
- [HPA 2010] Hart D, Wall BF, Hiller MC, Shrimpton PC. Frequency and collective dose for medical and dental X-ray examinations in the UK, 2008. Health Protection Agency. HPA-CECE-012. Chilton 2010.
- [ICRP 1991] The 1991 Recommendations of the International Commission on Radiological Protection. Publication 60. Annals of the ICRP, 21 (1-3) 1991.
- [ICRP 2007] The 2007 Recommendations of the International Commission on Radiological Protection. Publication 103. Annals of the ICRP, 37 (2-4) 2007.
- [IRSN-IVS 2010] Etard C, Sinno-Tellier S, Aubert B. Exposure of French Population by Ionizing Radiation due to Medical Diagnostic Examinations in 2007 (in French). Joint Report of the Institut de Radioprotection et de Sûreté Nucléaire and the Institut de Veille Sanitaire. Paris 2010.
- [Järvinen 2011] Järvinen H. Finnish Radiation and Nuclear Safety Authority (STUK). Personal Communication 2011.
- [Karthikesalingam et al. 2009] Karthikesalingam A, Markar S, Weerakkody R, Walsh SR, Carroll N, Praseedom RK. Radiation exposure during laparoscopic cholecystectomy with routine intraoperative cholangiography. Surgical endoscopy. 23(8):1845-1848; 2009
- [LRaP 2004] Swiss Federal Act on Radiation Protection (LRaP 814.50). Swiss Confederation. Bern; 2004
- [Ludlow et al. 2008] Ludlow J, Davies-Ludlow L, White SC. Patient risk related to common dental radiographic examinations: the impact of 2007 International Commission on Radiological Protection recommendations. The journal of the American Dental association. 139:1237-1243; 2008
- [Martin 2008] Martin CJ. Radiation dosimetry for diagnostic medical exposures. Radiation protection dosimetry. 128(4):389-412; 2008
- [Mettler et al. 2008] Mettler F, Huda W, Yoshizumi T, Mahesh M. Effective doses in radiology and diagnostic nuclear medicine: a catalog. Radiology. 248(1):254-63; 2008
- [Mini and Poretti 1984] Mini RL, Poretti G. Die Bestimmung der Strahlenbelastung einer Bevölkerungsgruppe gemäss ICRP26. Tagungsbericht der Schweizerischen Gesellschaft für Strahlenbiologie und Strahlenphysik; 1984

- [Mini 1992] Mini RL. Dosisbestimmungen in der medizinischen Röntgendiagnostik. Kerzers: Verlag Max Huber; 1992
- [NCRP 2009] National Council on Radiation Protection and Measurements. Ionizing Radiation Exposure of the Population of the United States. NCRP Report No. 160, Bethesda; 2009
- [NRPA 2009] A survey and inspection based on new radiation protection regulations. StrålevernRapport 2009:2. Østerås: Norwegian Radiation Protection Authority; 2009 Language: Norwegian
- [NRPA 2010] Radiology in Norway Anno 2008. Trends in examination frequency and collective effective dose to the population. StrålevernRapport 2010:12. Østerås: Norwegian Radiation Protection Authority; 2010 Language: Norwegian.
- [Pantos et al. 2009] Pantos I, Patatoukas G, Katritsis DG, Efstathopoulos E. Patient radiation doses in interventional cardiology procedures. Current cardiology reviews. 5(1):1-11; 2009
- [Perisinakis et al. 2003] Perisinakis K, Damilakis J, Grammatikakis J, Theocharopoulos N, Gourtsoyiannis N. Radiogenic risks from hysterosalpingography. European radiology. 13(7):1522-1528; 2003
- [Perisinakis et al. 2004] Perisinakis K, Damilakis J, Theocharopoulos N, Papadokostakis G, Hadjipavlou A, Gourtsoyiannis N. Patient exposure and associated radiation risks from fluoroscopically guided vertebroplasty or kyphoplasty. Radiology. 232(3):701-707; 2004
- [Poretti et al. 1971] Poretti G, Ionesco R, Lanz W. Erhebung über die Strahlenbelastung der Schweizer Bevölkerung infolge Röntgendiagnostischer Untersuchungen, Hrsg., Schweiz. Ver-einigung für Atomenergie; 1971
- [RIVM 2010] De Waard IR and Stoop P. Information System on Medical Radiation Applications: Data from the Reporting year 2008 (in Dutch). Rijksinstituut voor Volksgezondheid en Milieu. RIVM Report 300081005. Bilthoven; 2010.
- [Roser 2011] Roser H. Personal communication; 2011.
- [Ruiz Cruces et al. 1998] Ruiz Cruces R, García-Granados J, Diaz Romero F, Hernández Armas J. Estimation of effective dose in some digital angiographic and interventional procedures. The British journal of radiology. 71(841):42-7; 1998
- [Samara et al. 2010] Samara ET, Aroua A, Stauffer J-C, Bochud F, Verdun FR. Fluoroscopy guided procedures in cardiology: are we reducing patient exposure over time? Radiat Prot Dosimetry. 139(1-3):271-4; 2010
- [Samara et al. 2011] Samara ET, Aroua A, Trueb PhR, Stuessi A, Treier R, Schmidt S, Stauffer J-C, Bochud F, Zeller W, Verdun FR. An audit of diagnostic reference levels in interventional cardiology and radiology: are there differences between academic and non-academic centres? Radiat Prot Dosimetry (Epub Ahead of Print, PMID: 21335331).
- [Smith et al. 2009] Smith IR, Rivers JT, Hayes J, Stafford W, Codd C. Reassessment of radiation risks from electrophysiology procedures compared to coronary angiography. Heart Lung Circ. 18(3):191-199; 2009
- [Şorop et al. 2008] Şorop I, Mossang D, Iacob M, Dadulescu E, Iacob O. Update of diagnostic medical and dental x-ray exposures in Romania. Journal of radiological protection. 28(4):563-571; 2008
- [Tapiovaara et al. 1997] Tapiovaara M, Lakkisto M, Servomaa A. PCXMC—A PC-based Monte Carlo program for calculating patient doses in medical X-ray examinations (Helsinki: finnish centre for radiation and nuclear safety (STUK); 1997

- [Thomas et al. 2006] Thomas S, Kalkwarf H, Buckley D, Heubi JE. Effective dose of dual-energy X-ray absorptiometry scans in children as a function of age. *Journal of Clinical Densitometry*. 8(4):415-422; 2006
- [Treier et al. 2010] Treier R, Aroua A, Verdun FR, Samara E, Stuessi A, Trueb PR. Patient doses in CT examinations in Switzerland: implementation of national diagnostic reference levels. *Radiat Prot Dosimetry*. 142(2-4):244-54; 2010
- [Tsalafoutas et al. 2005] Tsalafoutas I, Chrysovergis D, Maniatis PN, Vassilaki AE, Triantopoulou CC, Yakoumakis EN, Kouliantianos ED. Radiation doses to patients from enteroclysis. *Radiation protection dosimetry*. 113(2):162-7; 2005
- [Frede Unhjem 2011] Unhjem F. Personal communication; 2011
- [UNSCEAR 2010] United Nations Scientific Committee on the Effects of Atomic Radiation. UNSCEAR 2008 Report Vol. I: Sources of Ionizing Radiation. United Nations. New York; 2010.
- [Yakoumakis et al. 2001] Yakoumakis E, Tsalafoutas I, Nikolaou D, Nazos I, Kouliantianos E, Proukakis C. Differences in effective dose estimation from dose-area product and entrance surface dose measurements in intravenous urography. *British Journal of Radiology*. 74(884):727-734; 2001
- [Zuppinger et al. 1961] Zuppinger A, Minder W, Sarasin R, Schär M. Die Strahlenbelastung der schweizerischen Bevölkerung durch röntgendiagnostische Massnahmen. *Radiol Clin* 30:1-5; 1961.

Appendices

Appendix 1: The “raddose.ch” website

Appendix 2: List of examination types according to EC Radiation Protection 154

Appendix 3: Dosimetry

Appendix 4: The dose vector

Appendix 5: Geographic distribution of the healthcare providers (total and respondents)

Appendix 6: Sensitivity analysis

Appendix 7: Examination frequency by healthcare providers

Appendix 8: Annual collective dose by healthcare providers

Appendix 9: Number of examinations (N), Collective dose (S), Frequency of examinations (N/1000 population), and average effective dose (E/caput)

Appendix 1:

The “raddose.ch” website

In der Schweiz stellt die durchschnittliche Strahlendosis der Bevölkerung durch medizinische Anwendungen den grössten Anteil der künstlichen Bestrahlungen dar. Er macht etwa ein Viertel der gesamten Strahlendosis der Schweizer Bevölkerung aus, welche 4 Millisievert (mSv) pro Jahr und Person beträgt.



DURCHSCHNITTliche STRAHLENDOSIS DER SCHWEIZER BEVÖLKERUNG
(pro Jahr und pro Person) - Quelle: Bundesamt für Gesundheit BAG

ERHEBUNG 2008

Aus Sicht der Globalen Initiative für Strahlenschutz in der Gesundheitsfürsorge der Weltgesundheitsorganisation „fordert die breite Nutzung von Strahlung in der Medizin einen Ansatz in der öffentlichen Gesundheit um die Gesundheitsrisiken zu kontrollieren bzw. zu minimieren bei gleichzeitiger Maximierung der Vorteile“. In diesem Kontext wird die Bewertung der Exposition der Bevölkerung der durch medizinische Anwendungen resultierenden Strahlungen mittels regelmäßiger nationaler Erhebungen gefördert.

Alle 5-10 Jahre sammelt und analysiert der **wissenschaftliche Ausschuss der Vereinten Nationen über die Effekte der atomaren Strahlung** (UNSCEAR) Daten auf globaler und regionaler Ebene zur Entwicklung der Exposition des Menschen bezüglich ionisierender Strahlung. Die periodischen UNSCEAR Erhebungen umfassen insbesondere die medizinische Exposition und liefern den globalen Trend der Häufigkeit der Untersuchungen. Die Ergebnisse der letzten Erhebung werden im Jahr 2009 veröffentlicht.

Nach Angaben der **Europäischen Leitlinien für die Einschätzung der Populationsdosis bei medizinischen Röntgenverfahren** (Europäische Kommission, Radiation Protection N°154, 2008), „wird empfohlen, die Häufigkeit der Erhebungen wenn möglich alle 5 Jahre zu wiederholen.“

Gemäss dem **Schweizer Strahlenschutzgesetz** (StSG), „fordert der Bund die wissenschaftliche Forschung über Strahlenwirkungen und Strahlenschutz.“ (Art. 5),

und „gibt der mit der medizinischen Untersuchung beauftragte Arzt der Aufsichtsbehörde die Daten bekannt, die für die medizinische Überwachung und das Erstellen von Statistiken notwendig sind. Die Aufsichtsbehörde darf diese Daten weder zu anderen Zwecken verwenden noch sie an Dritte weitergeben.“ (Art. 14)

Die Schweiz hat eine lange Tradition bezüglich Erhebungen über Exposition der Bevölkerung durch die Röntgendiagnostik, welche bis zum Ende der 50er Jahre zurück geht. Die letzte Updatestudie 2003 wurde vom **Bundesamt für Gesundheit** und vom **Schweizerischen Nationalfonds zur Förderung der wissenschaftlichen Forschung** unterstützt.

Die **Erhebung 2008** steht unter der Schirmherrschaft des Bundesamtes für Gesundheit (**BAG**) gemeinsam mit dem Universitätsinstitut für angewandte Strahlenphysik (**IRA**) und dem Universitätsinstitut für Sozial- und Präventivmedizin (**IUMSP**) und der Universitätsabteilung für Medizin und Öffentliche Gesundheit (**DUMSC**) der Universitätsklinik des Kantons Waadt (**CHUV**). Sie wird von einer **Begleitgruppe** mit Vertretern aus allen wissenschaftlichen Gesellschaften und Berufsverbänden begleitet.

RAD+DOSE EXPOSURE OF THE SWISS POPULATION BY MEDICAL RADIOLOGY

Home Kontext Erhebung 2008 Formular 2008 Ausfüllen Frühere Erhebungen Kontakt Deutsch | Français | Italiano | English

Erhebung 2008 PDF | Drucken | E-Mail

Erhebung zur Strahlenexposition der Bevölkerung durch die Röntgendiagnostik

Seit Ende der 50er-Jahre wird in der Schweiz die Strahlenexposition der Bevölkerung durch die Röntgendiagnostik regelmässig untersucht. Die letzte landesweite Erhebung bezog sich auf die Daten von 1998. Sie wurde unter der Federführung des Bundesamtes für Gesundheit (BAG), des Universitätsinstituts für angewandte Strahlenphysik (IRA), dem Universitätsinstitut für Sozial- und Präventivmedizin (IUMSP) und der Universitätsabteilung für Medizin und Öffentliche Gesundheit (DUMSC) der Universitätsklinik des Kantons Waadt gemeinsam durchgeführt. Diese Erhebung lieferte eine erhebliche Menge von Informationen zur Häufigkeit der in der Schweiz durchgeführten Untersuchungen und zu den damit verbundenen Dosen. Fünf Jahre danach wurde eine Aktualisierungserhebung anhand einer kleinen geschichteten Stichprobe durchgeführt. Damit konnte zum Beispiel festgestellt werden, dass der Einsatz der Computertomographie (CT) zwischen 1998 und 2003 stark zugenommen hat.

Aufgrund der raschen technologischen Entwicklung im Bereich der medizinischen Radiologie und der Weiterentwicklung der Praxis legen die internationalen Empfehlungen alle zehn Jahre eine Neubeurteilung der Situation nahe. Im Fall der Schweiz sollte somit die nationale Erhebung 2009-2010 anhand der Daten von 2008 erneut durchgeführt werden.

Bei den bisherigen Erhebungen wurden hauptsächlich Papierformulare verwendet, die an die Teilnehmerinnen und Teilnehmer versandt wurden. Für die Befragten war dies mit einem hohen Aufwand verbunden, der sich nur schwerlich wiederholen lässt. Daher wird bei dieser Erhebung die Möglichkeiten für eine Automatisierung der Datensammlung geprüft und genutzt. Vor allem sollen die Tarmed-Tariffcodes herangezogen werden.

Die europäische Arbeitsgruppe „Dose Datamed“ hat Empfehlungen für die Durchführung der nationalen Erhebungen zur Strahlenexposition der Bevölkerung erarbeitet. Diese Leitlinien, die von der Europäischen Kommission genehmigt wurden, werden bei dieser schweizerischen Erhebung berücksichtigt.

Mit der Erhebung wird die Entwicklung der Häufigkeit der Untersuchungen und der Dosen evaluiert. Für die Häufigkeiten werden alle Strukturen befragt, die medizinische Leistungen erbringen (Spitäler, Arztpraxen, Institute usw.). Einbezogen sind auch die Organisationen, die über die für die Studie relevanten Datenbanken verfügen (Santésuisse, kantonale Ärztesellschaften usw.). Für die Erhebung der Dosen werden die Daten der zahlreichen Studien, die in den letzten drei Jahren durchgeführt wurden, beigezogen. Bei Bedarf werden diese Daten durch gezielte Untersuchungen ergänzt. Bei dieser Gelegenheit kann die « Nationale Dosis Datenbank » aufgebaut werden, das in der Revisionsvorlage der Strahlenschutzverordnung vorgesehen ist.

Die Erhebung wird von einer Steuergruppe geleitet, in der das BAG, das IRA und das IUMSP vertreten sind, und von einer Begleitgruppe unterstützt, der Vertreterinnen und Vertreter aller betroffenen Fachgesellschaften und Berufsorganisationen angehören.

Begleitgruppe der Erhebung 2008

- Association Romande de Radioprotection
- Bundesamt für Gesundheit
- Die Spitäler der Schweiz
- Eidgenössische Kommission für Strahlenschutz und Überwachung der Radioaktivität
- Kollegium für Hausarztmedizin
- Santésuisse
- Schweizerische Chiropraktoren-Gesellschaft
- Schweizerische Gesellschaft für Allgemeinmedizin
- Schweizerische Gesellschaft für Gastroenterologie
- Schweizerische Gesellschaft für Innere Medizin
- Schweizerische Gesellschaft für Kardiologie
- Schweizerischen Gesellschaft für Pädiatrie
- Schweizerische Gesellschaft für Radiologie
- Schweizerische Gesellschaft für Strahlenbiologie und medizinische Physik
- Schweizerische Gesellschaft für Urologie
- Schweizerische Vereinigung der Fachleute für medizinisch-technische Radiologie
- Schweizerische Zahnärzte-Gesellschaft
- Vereinigung der Kantonsärztinnen und Kantonsärzte der Schweiz

Copyright © 2009 RAD+DOSE
All Rights Reserved

RAD+DOSE EXPOSURE OF THE SWISS POPULATION BY MEDICAL RADIOLOGY

Home Kontext Erhebung 2008 Formular 2008 Ausfüllen Frühere Erhebungen Kontakt Deutsch | Français | Italiano | English

Frühere Erhebungen PDF | Drucken | E-Mail

Berichte über die Erhebung 2003

#	Article Title
1	Die Strahlenexposition der Bevölkerung durch Röntgendiagnostik ist angestiegen Bundesamt für Gesundheit Bulletin 24/06:471-473, Bern 12. Juni 2006
2	Augmentation de l'exposition aux rayonnements dus au radiodiagnostic Bulletin de l'Office fédéral de la santé publique 24/06:471-473, Berne 12 juin 2006
3	Exposure of the Swiss population by radiodiagnosics: 2003 Review University Institute of Radiation Physics, Lausanne (2005) Aroua A, Vallet JP, Verdun FR and Vader JP

Artikel über die Erhebung 2003

#	Article Title
1	Exposure of the Swiss population by radiodiagnosics: 2003 review <i>(* beschränkter Zugriff)</i> Health Phys 92(5):442-8 (2007) Aroua A, Vader JP, Vallet JF and Verdun FR

Berichte über die Erhebung 1998

#	Article Title
1	Enquête sur l'exposition par le radiodiagnostic en Suisse en 1998 Institut Universitaire de Radiophysique, Lausanne 2000 A.Aroua A, Vader JP and Vallet JF
2	A survey on exposure by Radiodiagnosics in Switzerland in 1998 University Institute of Radiation Physics, Lausanne (2000) Aroua A, Vader JP and Vallet JF

Artikel über die Erhebung 1998

#	Article Title
1	Number of x-ray examinations performed on paediatric and geriatric patients compared with adult patients <i>(* beschränkter Zugriff)</i> Radiat Prof Dosim 123(3):402-8 (2007) Aroua A, Bochud FO, Vallet JF, Vader JP and Verdun FR
2	CT utilisation in Switzerland: frequency and secular trends Journal de Radiologie 87(9):1057-62 (2006) Mildez B, Aroua A, Verdun FR, Vallet JF, Schnyder P and Vader JP
3	Radiation Exposure in Dental Radiology: A 1998 Nation-wide Survey in Switzerland <i>(* beschränkter Zugriff)</i> Dentomaxillofacial Radiology 33:211-219 (2004) Aroua A, Buchillier-Decker I, Dula K, Nedjadi Y, Perrier M, Vader JP and Vallet JF

Info Rad+Dose

Adresse:

Institut Universitaire de Radiophysique Appliquée
 Enquête 2008 sur l'exposition par le radiodiagnostic
 Rue du Grand-Pré 1
 CH - 1007 Lausanne



E-Mail: survey2008@raddose.ch

Telefon: + 41 (0)21 31 48 068

Fax: + 41 (0)21 31 48 299

Informationen herunterladen als vCard

[*] Ihr Name

[*] E-Mail-Adresse

[*] Betreff

[*] Ihre Nachricht

Eine Kopie dieser Nachricht an Ihre E-Mail-Adresse senden



Geben Sie den angezeigten Sicherheitscode ein.

Senden

ONLINE-FORMULAR

Bitte melden Sie sich an, um auf den internen Bereich der Website zugreifen zu können!

Benutzername

Passwort

Angemeldet bleiben

Anmelden

Passwort vergessen?



PAPIERFORM

Wenn Sie möchten, können Sie das Formular in Papierform ausfüllen. Klicken Sie auf dem unten stehenden Link um die Datei herunterzuladen:

Herunterladen (Druckbares Format)



Sie benötigen das Programm Adobe® Reader® oder ähnliches um die Datei anzeigen bzw. drucken zu können:



RAD+DOSE EXPOSURE OF THE SWISS POPULATION BY MEDICAL RADIOLOGY

Home Kontext Erhebung 2008 Formular 2008 Ausfüllen Frühere Erhebungen Kontakt Sich abmelden Deutsch | Français | Italiano | English

[Profil](#) >> [Anlagenliste](#) >> [Formular](#) >> [Die Daten validieren](#)

Ihre Angaben ändern

Wichtig:

- Bitte ergänzen Sie die Informationen Ihrem Benutzerprofil.
- Pflichtfelder müssen ausgefüllt werden, um den Zugriff auf das Formular zu ermöglichen.
- Nach dem Update Ihrem Profil, **aktualisieren Sie die Anlagenliste** >>

Username: VD832B1

Benutzer: CHUV
Dépt. Radiologie médicale
Radiodiag. et radiol. inter.

Medizinisches Fachgebiet: Radiologie

Adresse: Rue du Bugnon 46 BH-10-117
1011 Lausanne

Kontakt & Bearbeitung: Geben Sie bitte die Daten des Ansprechpartners ein.

Vorname: Stéphane

Nachname: Coendoz

Position: Directeur administratif

Telefon: 021.314.45.67

Fax:

Email-Adresse: Stephane.Coendoz@chuv.ch

Anmerkungen:

Statut Contribution:

Pflichtfeld Information: Bewegen Sie die Maus auf das Icon

Copyright © 2009 RAD+DOSE
All Rights Reserved

RAD+DOSE EXPOSURE OF THE SWISS POPULATION BY MEDICAL RADIOLOGY

Home Kontext Erhebung 2008 Formular 2008 Ausfüllen Frühere Erhebungen Kontakt Sich abmelden Deutsch | Français | Italiano | English

[Profil](#) >> [Anlagenliste](#) >> [Formular](#) >> [Die Daten validieren](#)

ANLAGENLISTE

Zeichenerklärung

Kürzel	Anwendung
BES	Beschleuniger
CT	Computer Tomographie
DVT	Digitale Volumentomographie (dental)
FR	Fernröntgenaufnahme (dental)
KDM	Knochendichtemessung
MAM	Mammographie
OPF	Panorama + Fernröntgenaufnahme (dental)
OPG	Panorama-Schichtaufnahme
RA	Radiographie (nur)
RAD	Radiographie + Fluoroskopie
RD	Fluoroskopie (nur)
RDI	Fluoroskopie interventionell (intensive Dosis)
RKL	Kleine Dental Röntgenanlage (< 70 kV)

- Bitte bezeichnen Sie die Anlagen, die die eingegebene Daten **mit eingeschlossen** sind.
- Die Schaltfläche **Edit** anklicken um die Einstellung zu ändern.
- Um **eine neue Anlage einzufügen** klicken Sie **New Unit** an.

Anlage-ID	Anwendung	Kurzbezeichnung	mit eingeschlossen	
VD-832-1-17	RA	MAXIMUS CM80	✓	Edit
VD-832-1-24	RKL	RITTER D502	✓	Edit
VD-832-1-31	RKL	ORALIX 65	✓	Edit
VD-832-1-32	RKL	ORALIX 65	✓	Edit
VD-832-1-34	RKL	ORALIX AC	✓	Edit
VD-832-1-38	RAD	SIREMOBIL 2000	✓	Edit
VD-832-1-42	RA	MPH 80	✓	Edit
VD-832-1-43	RDI	ADVANTIX VMP 100	✓	Edit
VD-832-1-45	RAD	ADVANTX	✓	Edit
VD-832-1-46	RDI	ADVANTX	✓	Edit
VD-832-1-48	RD	SIREMOBIL COMPACT	✓	Edit
VD-832-1-49	RD	SIREMOBIL ISO C	✓	Edit
VD-832-1-50	RD	SIREMOBIL COMPACT	✓	Edit
VD-832-1-54	OPF	ORTHOPHOS XG PLUS	✓	Edit
VD-832-1-56	RAD	ADVANTIX VMP 100	✓	Edit
VD-832-1-57	RD	BV 312	✓	Edit
VD-832-1-59	RAD	SIREMOBIL COMPACT	✓	Edit
VD-832-1-60	CT	CT LIGHTSPEED ULTRA ADVANTAGE	✓	Edit
VD-832-1-61	RA	DR 5100	✓	Edit
VD-832-1-65	CT	CT LIGHTSPEED VCT	✓	Edit
VD-832-1-67	RDI	ALLURA FD10/20 BIPLAN	✓	Edit
VD-832-1-68	MAM	CONTOUR 2000	✓	Edit
VD-832-1-69	MAM	LORAD MULTI-CARE	✓	Edit
VD-832-1-70	RA	OPTIMUS 80	✓	Edit
VD-832-1-71	RA	SENOGRAPH ESSENTIAL	✓	Edit
VD-832-1-72	RDI	INTEGRIS ALLURA FD20C	✓	Edit
VD-832-1-73	RD	AXIOM ARTIS DFC	✓	Edit
VD-832-1-74	RA	OPTIMUS 65	✓	Edit
VD-832-1-75	RA	OPTIMUS 65	✓	Edit
VD-832-1-9	RA	MAXIMUS CM80	✗	Edit

Copyright © 2009 RAD+DOSE

Profil » Anlagenliste » Formular » Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE | FLUOROSKOPIE | ANGIOGRAPHIE | CT | INTERVENTIONELLE | DENTAL | MAMMOGRAPHIE | KDM | **BEARBEITEN**

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Kopf	Orbitae	
	Schläfenbein (Felsenbein & Mastoid)	
	Keilbein (Türkensattel & Fissura sphenoidalis)	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Schädel" ein	1347
	Nasenbein	
	Sinus	
	Jochbogen	
	Temporomandibulargelenk (Kiefergelenk)	
	Hals-Hinterkopf-Winkelgelenk	
	Oberkiefer	
	Unterkiefer	
	Cephalometrie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Gesichtsknochen" ein	
	Dacryocystographie (Tränenkanäle)	
	Sialographie (Speicheldrüsen)	
Augen		
Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Kopf - Weichgewebe" ein	0	
Hals	HWS	1975
	Larynx	
	Rachen	
	Lufttröhre	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Hals - Weichgewebe" ein	0
Thorax	BWS	1330
	Schulterblatt(-blätter)	
	Schlüsselbein(e)	
	Akromioklavikulargelenk	
	Sternoklavikulargelenk	
	Manubriosternalgelenk	
	Sternum	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Schultergürtel" ein	4370
	Rippenthorax	
	Lunge	44779
Obere Thoraxapertur	0	
Bronchographie	0	

	Bronchographie	0	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Thorax" ein		
abdominal Region	LWS	3643	
	Lumbosakralgelenk	0	
	Abdomen	3690	
Becken Region	Os ilium / Os ischium / Os pubis		
	Sacrum		
	Iliosakralgelenk		
	Steißbein		
	Pelvimetrie (Geburtshilfe)		
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Beckenknochen" ein	5712	
	Hüftgelenk (ein oder beides)	4711	
	Pelvis (Weichgewebe)	0	
Extremitäten	Oberarm (Humerus)	620	
	Ellbogen	1652	
	Unterarm (Speiche & Elle)	817	
	Handgelenk (Os scapuloideum)	2787	
	Hand	2171	
	Finger	1022	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Unterarm, Handgelenk & Hand" ein		
	Oberschenkelknochen (Femur)	1082	
	Kniegelenk		
	Kniescheibe (Patella)		
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Knie" ein	6126	
	Unterschenkelknochen (Tibia & Fibula)	1661	
	Oberes Sprunggelenk (OSG)	4169	
	Fuss	4171	
	Kalkaneus (Fersenbein)	445	
	Zehen	761	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Unterschenkel, OSG & Fuss" ein		
		Ganze Schenkel	985
	Rumpf	Scoliosis (ganze Wirbelsäule)	541
	Kopf & Rumpf	Ganze Skelett (Skeletal survey)	0
Weitere			
Weitere			
Weitere			
Weitere			
Weitere			
Kommentar	Pour les examens de la région de la tête, les items proposés ne correspondent ni des actes statistiques utilisés dans notre département, ni à la nomenclature Tarmed.		
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Radiographie" ein		

Copyright © 2009 RAD+DOSE
All Rights Reserved

Profil » Anlagenliste » Formular » Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE FLUOROSKOPIE ANGIOGRAPHIE CT INTERVENTIONELLE DENTAL MAMMOGRAPHIE KDM **BEARBEITEN**

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Verdauungstrakt (Hals + Thorax + Abdomen)	Speiseröhre (Bariumschluck)	229
	Magen & Duodenum (Bariumbrei)	474
	Dünndarm (Dünndarmpassage)	0
	Enteroklysma (Dünndarmeinlauf)	33
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Speiseröhre & Magen & Dünndarm" ein	
	Kolon (Kolonkontrasteinlauf)	85
	Defäkographie	58
Gallengang	Retrograde Cholangiographie	0
	Intraoperative Cholangiographie	
	Intravenöse Cholangiographie	1
	T-Drain Cholangiographie	86
	Transhepatische Cholangiographie	
	Endoskopische retrograde Cholangiopankreatographie (ERCP)	
	Retrograde Pankreatographie	
	Cholecystographie	0
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Gallengang" ein	
Harn- und Geschlechtsapparat	Intravenöse Urographie	11
	Retrograde Pyelographie	1
	Nephrostographie	0
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Nieren & Harnleitern" ein	
	Retrograde Cystographie	0
	Miktions-Cysto-Urethrographie	314
	Urethrographie	0
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Blase & Harnröhre" ein	
Hysterosalpingographie (Frauenheilkunde)	107	
Rückenmark	Cervicale Myelographie	
	Thorakale Myelographie	
	Lumbale Myelographie	
	Sacrale Myelographie	
	Myelographie des ganzen Rückenmarks	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Myelographie" ein	47
Gelenke	Temporomandibulargelenk Arthrographie	
	Schulter Arthrographie	

Profil » Anlagenliste » **Formular** » Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE FLUOROSKOPIE **ANGIOGRAPHIE** CT INTERVENTIONELLE DENTAL MAMMOGRAPHIE KDM BEARBEITEN

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Blutgefäße	Cerebraliangiographie (Arteriographie)	73
	Orbitale Phlebographie	4
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Cerebrale Angiographie" ein	
	Koronarangiographie (Koronarographie)	1990
	Thorakale Aortographie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Herzangiographie (Angiokardiographie)" ein	
	Bronchialarteriographie	
	Pulmonalisarteriographie	9
	Obere Cavographie	560
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Thorakale Angiographie" ein	
	Abdominale Aortographie	46
	Nierenarteriographie	
	Mesenterialarteriographie	
	Untere Cavographie	150
	Nierenphlebographie	
	Nebennierenphlebographie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Abdominale Angiographie" ein	
	Beckenarteriographie	
	Ovarialphlebographie	
	Spermataphlebographie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Becken Angiographie" ein	
	Arteriographie der oberen Extremität	
	Phlebographie der oberen Extremität	189
Arteriographie der unteren Extremität	359	
Phlebographie der unteren Extremität	6	
Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Periphere Angiographie" ein		
Lymphgefäße	Thorakale Lymphangiographie	
	Abdominale Lymphangiographie	
	Beckenlymphangiographie	
	Lymphangiographie der oberen Extremität	
	Lymphangiographie der unteren Extremität	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Lymphangiographie" ein	

Profil » Anlagenliste » Formular » Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE FLUOROSKOPIE ANGIOGRAPHIE **CT** INTERVENTIONELLE DENTAL MAMMOGRAPHIE KDM BEARBEITEN

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Kopf	Orbitae	
	Schläfenbein (Felsenbein)	
	Temporomandibulargelenk (Kiefergelenk)	
	Sella turcica	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "CT Schädel" ein	
	Gesicht	1092
	Dentale (Zähne)	13
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "CT Gesichtsknochen" ein	
	Gehirn	
	Fossa posterior	
	Gehirn vaskulär	
	Hypophyse	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "CT Gehirn" ein	7207
	Sinus	
Innerer Gehörgang		
Nasenhöhle		
Mund		
Total "CT Kopf - Weichgewebe" ein		
Hals	HWS	
	Hals	
	Larynx	
	Rachen	
	Hals vaskulär	
Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "CT Hals" ein	2427	
Thorax	BWS	
	Mediastinum	
	Lunge standard	
	Lunge hohe Auflösung	
	Herz	
	Thorakale Aorta	
	Lunge vaskulär	
Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "CT Thorax" ein	8324	
Abdomen	LWS	
	Ganzes Abdomen	8303

Profil » Anlagenliste » Formular » Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE | FLUOROSKOPIE | ANGIOGRAPHIE | CT | **INTERVENTIONELLE** | DENTAL | MAMMOGRAPHIE | KDM | BEARBEITEN

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Kopf & Hals	Zerebrale Dilatation/Stenting	
	Zerebrale Embolisation (AVM, Aneurysma, Tumor)	
	Zerebrale Thrombolyse	
	Kopf & Hals Punktion	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Zerebrale Interventionen" ein	55
Thorax	Koronardilatation/Stenting (PTCA)	916
	Pacemaker Implantation (temporär oder permanent)	15
	Zentraler Venenkatheter Durchführung (Hickman line)	
	Herz Thermoablation	
	Valvuloplastie	23
	Inferior Vena Cava (IVC) Filter Fitting	88
	Oesophagusdilatation/Stenting	
	Thorakal Dilatation/Stenting	
	Thorakal Embolisation	
	Thorakal Thrombolyse	
	Thoraxbiopsie	
	Electrophysiologie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Andere Thoraxinterventionen" ein	
Abdomen	Gallengangserweiterung/Stenting	
	Gallengangdrainage	
	Steinextraktion aus dem Gallengang	
	Nierenarteriendilatation/Stenting	
	Nierendrainage	
	Lithotripsie	
	Nephrostomie	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Gallengangssystem & Harnsystem" ein	
	TIPS (Leber)	
	Abdominale Dilatation/Stenting	
	Abdominale Embolisation	
	Abdominale Thrombolyse	
	Abdominael Punktion	
Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Abdominale Interventionen" ein		
Becken	Beckengefäßdilatation	
	Beckengefäßembolisation	

Profil >> Anlagenliste >> **Formular** >> Die Daten validieren

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE FLUOROSKOPIE ANGIOGRAPHIE CT INTERVENTIONELLE **DENTAL** MAMMOGRAPHIE KDM BEARBEITEN

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Zähne & Zahnfleisch	Apikal (1-2 Filme)	
	Apikal (mehr als 2 Filme)	
	Apikal (Status)	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl "Apikal" ein	
	Kurze Bissflügelaufnahme (1-2 Filme)	
	Lange Bissflügelaufnahme (mehr als 2 Filme)	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Bissflügelaufnahmen" ein	
	Aufbissaufnahme	
	Panorama-Schichtaufnahme (OPG)	1169
	CT (Unterkiefer/Oberkiefer)	13
Kopf	Schädel	
	Fernröntgenaufnahme	
	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl des Bereichs "Kopf" ein	
Extremitäten	Hand	
Weitere		
Weitere		
Weitere		
Kommentar	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Dentalradiologie" ein	

[Profil](#) >>> [Anlagenliste](#) >>> [Formular](#) >>> [Die Daten validieren](#)

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE | FLUOROSKOPIE | ANGIOGRAPHIE | CT | INTERVENTIONELLE | DENTAL | **MAMMOGRAPHIE** | KDM | **BEARBEITEN**

	Untersuchungstyp	Anzahl Untersuchungen im 2008
Brust	Symptomatisch (1-2 Aufnahmen / 1 Brust oder beide Brüste)	1937
	Screening (1-2 Aufnahmen / beide Brüste)	1058
Kommentar	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Mammographie" ein	

[Profil](#) >>> [Anlagenliste](#) >>> [Formular](#) >>> [Die Daten validieren](#)

ERHEBUNG 2008 ÜBER DIE EXPOSITION DURCH DIE RÖNTGENDIAGNOSTIK (FORMULAR)

RADIOGRAPHIE | FLUOROSKOPIE | ANGIOGRAPHIE | CT | INTERVENTIONELLE | DENTAL | **MAMMOGRAPHIE** | **KDM** | **BEARBEITEN**

	Untersuchungstyp	Anzahl Untersuchungen im 2008
abdominal Region	LWS	
Becken Region	Hüftgelenk	
Extremitäten	Obere / Untere Extremität	
Weitere	Densitométrie toute région	1173
Kommentar	Wenn die detaillierte Zahlen nicht verfügbar sind, geben Sie die Gesamtzahl der "Knochendichtemessung" ein	

[Profil](#) >>> [Anlagenliste](#) >>> [Formular](#) >>> [Die Daten validieren](#)

DIE DATEN VALIDIEREN

Validieren Sie Ihre Eingegebenen Daten.

Nach dem Validieren, können die Daten des Formulars **nicht geändert werden** und werden berücksichtigt für die statistische Bearbeitung.

Um das Formular wieder zu aktivieren, wird nötig sein den Administrator zu kontaktieren.

Validate

Appendix 2:

List of examination types
according to EC Radiation
Protection 154

Table 1: Plain film radiography

Region of body	Specific exam types	DOSE DATAMED exam categories
Head	Skull <ul style="list-style-type: none"> - Orbits - Temporal bones <ul style="list-style-type: none"> - petrous bone - mastoids - Sphenoid bone <ul style="list-style-type: none"> - sella turcica - sphenoid fissures Facial bones <ul style="list-style-type: none"> - Nose - Sinuses - Zygomax - Temporo-mandibular joint - Cervico-occipital hinge - Maxilla - Mandible - Cephalometry 	Skull & facial bones
	Dacryocystography (tear ducts) Sialography (salivary glands) Eyes/orbits	Head - soft tissue
	Neck	Cervical spine Larynx Pharynx Trachea
Chest/Thorax	Thoracic spine	Thoracic spine
	Shoulder blades/ scapulae Collar bone(s) / clavicle(s) Acromio-clavicular joint Sterno-clavicular joint Manubrio-sternal joint Sternum	Shoulder girdle
	Ribs	Ribs
	Lung Thoracic inlet Bronchography	Chest/thorax/lung
Abdomen	Lumbar spine	Lumbar spine
	Lumbo-sacral joint	Lumbo-sacral joint only
	Abdomen (plain film, patient supine or erect)	Abdomen

Pelvis	Pelvic bones - Ilium/ischium/pubis - Sacrum - Sacro-iliac joint - Coccyx	Pelvic bone
	Pelvimetry (obstetric)	
	1 or both hips	Hips
	Pelvis (soft tissue)	Pelvis (soft tissue)
Limbs	Upper arm (humerus)	Upper arm
	Elbow	Elbow
	Forearm (radius & ulna) Wrist (scaphoid) Hand - Fingers & thumbs	Forearm, wrist & hand
	Femur	Femur
	Knee Knee cap (patella)	Knee
	Lower leg (tibia & fibula) Ankle Foot Calcaneum (heel) Toes	Lower leg, ankle & foot
	Whole leg	Leg length
Trunk	Scoliosis	Whole spine
Head & trunk	Whole skeleton	Skeletal survey
Teeth & gums	1-2 periapical films 1-2 bitewing films 1 occlusal film	Intra-oral <3 films
	>2 periapical films Periapical full mouth survey >2 bitewing films	Intra-oral >2 films
	Panoramic full mouth scan	Panoramic
Breast	Symptomatic: - 1 or 2 views of 1 or both breasts Screening: - 1 or 2 views of both breasts	Mammography

Table 2: Radiography/fluoroscopy (excluding interventional procedures)

Region of body	Specific exam types	DOSE DATAMED exam categories
GI tract (Neck + chest + abdomen)	Oesophagus (Ba swallow) Stomach & duodenum (Ba meal) Small intestine (Ba follow) Enteroclysis (small intestine enema)	Oesoph. & stomach & small intestine
	Colon (Ba enema)	Colon
	Defecography	Defecography
Biliary tract	Retrograde cholangiography Operative cholangiography Intravenous cholangiography T drain cholangiography Transhepatic cholangiography Endoscopic retrograde cholangio-pancreatography (ERCP) Retrograde pancreatography Cholecystography	Biliary tract
Uro-genital tract	Intravenous urography (IVU)	IVU
	Retrograde pyelography Nephrostography	Kidneys & ureters
	Retrograde cystography Micturitional cysto-urethrography (MCU) Urethrography	Bladder & urethra
	Hysterosalpingography	Gynaecological
Spinal cord	Cervical myelography Thoracic myelography Lumbar myelography Sacral myelography Whole spine myelography	Myelography
Joints	Temporal-mandibular joint arthrography Shoulder arthrography Hip arthrography Elbow arthrography Wrist arthrography Knee arthrography Ankle arthrography	Arthrography

Angiography	Cerebral angiography Petrous phlebography	Cerebral angiography
	Coronary angiography (CA) - coronary arteries only - cor arts + L ventricle - cor arts + L ventricle + aorta Thoracic aortography	Cardiac angiography (angiocardiography)
	Bronchial arteriography Pulmonary arteriography Upper venacavography	Thoracic angiography
	Abdominal aortography Renal arteriography Mesenteric arteriography Lower venacavography Renal phlebography Suprarenal phlebography	Abdominal angiography
	Pelvic arteriography Ovarian phlebography Spermatic phlebography	Pelvic angiography
	Upper & lower limb arteriography Upper & lower limb phlebography	Peripheral angiography
Lymphangiography	Thoracic lymphangiography Abdominal lymphangiography Pelvic lymphangiography Upper & lower limb lymphangiography	Lymphangiography

Table 3: CT Examinations

Region of body	Specific exam types	DOSE DATAMED exam categories
Head	Skull <ul style="list-style-type: none"> - Orbits - Temporal bone <ul style="list-style-type: none"> - Petrous bone - Temporal-mandibular joint - Sella turcica Face Dental	Skull & facial bones
	Brain <ul style="list-style-type: none"> - Cerebrum - Posterior fossa - Brain vascular Pituitary gland	Brain
	Sinuses Internal auditory meatus Nasal cavity Mouth	Head soft tissues
Neck	Cervical spine	Cervical spine
	Neck Larynx Pharynx Neck vascular	Neck
Chest	Thoracic spine	Thoracic spine
	Mediastinum Lungs standard Lungs High Resolution Heart Thoracic aorta Lungs vascular	Chest/thorax
Abdomen	Lumbar spine	Lumbar spine
	Full abdomen Upper abdomen	Abdomen
	Liver / pancreas Kidneys / Supra-renal glands	Liver, pancreas & kidneys
	Hip / pelvic bone Sacrum/coccyx Sacro-iliac joint	Pelvic bones
Pelvis	Pelvimetry (obstetric)	Pelvimetry
	Pelvis (soft tissues/vascular)	Pelvis

Neck + chest + abdomen	Full spine	Full spine
Chest + abdomen	Chest/abdomen	Chest & abdomen
Abdomen + Pelvis	Abdomen/pelvis	Abdomen & pelvis
Chest + abdomen + pelvis	Whole trunk	Chest, abdomen & pelvis
Limbs	Shoulder Elbow Wrist Hand Leg Thigh Knee Calcaneum Ankle Foot	Limbs

Table 4: Interventional Radiology

Region of the body	Specific procedure types	DOSE DATAMED procedure categories
Head & neck	Cerebral dilatation/stenting Cerebral embolisation (AVM, aneurysm, tumor) Cerebral thrombolysis Head & neck puncture	Cerebral interventions
Chest	Coronary dilatation/stenting (PTCA)	PTCA
	Cardiac pacemaker fitting (temporary or permanent)	Pacemaker
	Central venous line fitting	Hickman line
	Cardiac thermo-ablation Valvuloplasty IVC (caval) filter fitting Oesophagus dilatation/stenting Thoracic dilatation/stenting Thoracic embolisation Thoracic thrombolysis Thoracic region biopsy Electrophysiology	Other thoracic intervents.
Abdomen	Bile duct dilatation/stenting Bile duct drainage Bile duct stone extraction Renal artery dilatation/stenting Renal drainage Lithotripsy Nephrostomy	Biliary & urinary systems
	TIPS (liver)	TIPS
	Abdominal dilatation/stenting Abdominal embolisation Abdominal thrombolysis Abdominal region biopsy	Abdominal interventions
Pelvis	Pelvic vessel dilatation Pelvic vessel embolisation Pelvic vessel thrombolysis	Pelvic interventions
Limbs	Upper limb dilatation Upper limb embolisation Upper limb thrombolysis Popliteal dilatation (behind knee) Lower limb dilatation Lower limb embolisation Lower limb thrombolysis Limbs biopsy	Limb interventions

Appendix 3: Dosimetry

Table 1: Technical parameters for radiography examinations

Examination	Projection	Tension (kV _p)	Tube filtration (mm Al)	SDD* (cm)	SSD** (cm)	Field size (at the detector level)	Current (mAs)	Remarks
Skull	Face /Lateral	70	3	120	100	20/25	50	Rare in practice
Face - Sinus	Face	70	3	120	100	18/24	64	
Neck – soft tissue	Lateral	63	3	120	90	18/24	32	
Cervical spine	Face/ Lateral	66	3	150	120	18/24	40	
Thoracic spine	Face	77	3	120	90	18/43	40	
Thoracic spine	Lateral	81	3 + filter comp	120	80	18/43	32	
Clavicle	Face/Neer	66	3	120	90	18/24	25	
Chest	Face	125	3	200	170	35/43	3	
Chest	Lateral	125	3	150	120	35/43	4	
Lumbar spine	Face	77	3	120	80	35/43	80	
Lumbar spine	Lateral	81	3	120	75	18/43	120	
Abdomen	Face	73	3	120	80	35/43	50	
Pelvis	Face	73	3	120	80	35/43	50	
Pelvic bones	Face	77	3	120	80	24/30	50	Direct exposition of the gonads
Hip	Axial/Lesquene	81	3	120	80	24/30	100	Direct exposition of the gonads
Forearm - Wrist - Hand	Face/Lateral /Oblique	48	3	120	115	18/24	10	
Femur	Face/ Lateral	70	3	120	100	18/43	40	
Knee	Face / Lateral	48	3	120	110	18/24	16	
Leg - Ankle - Foot	Face / Lateral	48	3	120	115	15/24	20	
Scoliosis (entire spine)	Face	70	3 + filter comp	200	170	35/100	40	

*SDD: Source-to-detector distance, **SSD: Source-to-skin distance

Table 2: Computed tomography dose index (CTDI) and dose-length product (DLP) for CT examinations

Protocol	Mean CTDI _{vol}	Median CTDI _{vol}	Mean DLP	Median DLP
Skull/brain	62	58	871	867
Brain (vascular)	56	57	847	860
Sinus	20	18	282	252
Petrous bone	56	50	509	370
Cervical spine	22	19	459	459
Neck	18	15	456	462
Shoulder	28	20	458	386
Thorax	10	10	330	338
Thorax (vascular)	12	12	370	353
Thorax/upper abdomen	12	13	494	449
Upper abdomen	13	12	384	367
Upper abdomen (vascular)	12	12	432	411
Abdomen/pelvis	12	11	507	497
Abdomen/pelvis (vascular)	15	13	620	607
Pelvis	17	15	429	381
Pelvis (vascular)	16	12	445	429
Thorax/abdomen/pelvis	14	13	734	702
Lumbar spine	28	23	592	494
Lower limbs	13	13	1049	695
Heart (cardiovascular)	49	39	814	763
Heart	14	8	679	122

Appendix 4: The dose vector

Table 1: Effective doses for radiography examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀	E 2008 ICRP₁₀₃
[Head] Orbits	4,10E-02	3,00E-02	4,00E-02
[Head] Temporal bones	1,07E-01	6,00E-02	8,00E-02
[Head] Sphenoid bone	1,05E-01	3,00E-02	4,00E-02
[Head] Nose	3,25E-03	4,00E-02	5,00E-02
[Head] Sinuses	1,12E-01	4,60E-02	5,75E-02
[Head] Zygomas	1,03E-01	4,00E-02	5,00E-02
[Head] Temporo-mandibular joint	1,66E-02	4,00E-02	5,00E-02
[Head] Cervico-occipital hinge	3,70E-02	4,20E-02	5,25E-02
[Head] Maxilla	3,70E-02	4,00E-02	5,00E-02
[Head] Mandible	3,70E-02	4,00E-02	5,00E-02
[Head] Cephalometry	3,70E-02	4,00E-02	5,00E-02
[Head] Dacryocystography	1,07E-01	8,00E-02	1,00E-01
[Head] Sialography	2,95E-01	4,00E-02	5,00E-02
[Head] Eyes	3,65E-02	4,00E-02	5,00E-02
[Neck] Cervical spine	5,16E-01	1,02E-01	1,07E-01
[Neck] Larynx	1,70E-02	5,00E-02	6,00E-02
[Neck] Pharynx	1,70E-02	5,00E-02	6,00E-02
[Neck] Trachea	7,24E-02	1,00E-01	1,20E-01
[Chest / Thorax] Thoracic spine	2,92E+00	4,35E-01	5,28E-01
[Chest / Thorax] Shoulder blades / scapulae	1,42E-01	2,90E-02	4,20E-02
[Chest / Thorax] Collar bone(s) / clavicle(s)	7,13E-03	2,90E-02	4,20E-02
[Chest / Thorax] Acromio-clavicular joint	5,85E-02	2,90E-02	4,20E-02
[Chest / Thorax] Sterno-clavicular joint	6,58E-02	2,90E-02	4,20E-02
[Chest / Thorax] Manubrio-sternal joint	3,73E-01	2,90E-02	4,20E-02
[Chest / Thorax] Sternum	3,02E-01	2,90E-02	4,20E-02
[Chest / Thorax] Ribs	2,18E-01	5,00E-02	6,00E-02
[Chest / Thorax] Lung	5,74E-02	5,05E-02	5,40E-02
[Chest / Thorax] Thoracic inlet	5,74E-02	3,80E-02	3,90E-02
[Chest / Thorax] Bronchography	5,74E-02	3,80E-02	3,90E-02
[Abdomen] Lumbar spine	3,44E+00	1,70E+00	1,50E+00
[Abdomen] Lumbo-sacral joint	1,27E+00	8,22E-01	4,58E-01
[Abdomen] Abdomen (plain film, patient supine or erect)	1,34E+00	7,80E-01	7,50E-01
[Pelvis] Ilium / ischium / pubis	7,24E-01	8,10E-01	6,75E-01
[Pelvis] Sacrum	1,95E+00	1,08E+00	9,00E-01
[Pelvis] Sacro-iliac joint	1,27E+00	1,08E+00	9,00E-01
[Pelvis] Coccyx	1,77E+00	1,08E+00	9,00E-01
[Pelvis] Pelvimetry (obstetric)	1,05E+00	5,40E-01	4,50E-01
[Pelvis] Hips (one or both)	1,24E+00	1,13E+00	6,30E-01
[Pelvis] Pelvis (soft tissue)	1,05E+00	5,40E-01	4,50E-01
[Limbs] Upper arm (humerus)	7,05E-02	6,10E-03	
[Limbs] Elbow	5,83E-04	6,00E-04	
[Limbs] Forearm (radius & ulna)	1,05E-03	1,10E-03	

[Limbs] Wrist (scaphoid)	2,48E-04	2,00E-04	
[Limbs] Hand	2,16E-04	2,00E-04	
[Limbs] Fingers & thumbs	7,70E-05	1,00E-04	
[Limbs] Femur	1,36E-02	4,00E-02	2,00E-02
[Limbs] Knee joint	4,47E-03	8,00E-04	4,00E-04
[Limbs] Knee cap (patella)	6,31E-04	4,00E-04	2,00E-04
[Limbs] Lower leg (tibia & fibula)	4,35E-03	4,00E-04	4,00E-04
[Limbs] Ankle	3,67E-04	4,00E-04	4,00E-04
[Limbs] Foot	4,92E-04	4,00E-04	4,00E-04
[Limbs] Calcaneum (heel)	3,39E-04	4,00E-04	4,00E-04
[Limbs] Toes	4,50E-05	4,00E-04	4,00E-04
[Limbs] Whole leg (Leg length)	1,36E-02	2,00E-02	2,00E-02
[Trunk] Scoliosis (Whole spine)	3,44E+00	3,40E-01	3,40E-01
[Head & trunk] Whole skeleton (Skeletal survey)	3,44E+00	3,44E+00	

Table 2: Effective doses for fluoroscopy examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀
[GI tract] Oesophagus	6,59E+00	1,10E+00
[GI tract] Stomach & duodenum	2,28E+01	1,20E+01
[GI tract] Small intestine	3,83E+01	3,80E+00
[GI tract] Enteroclysis	6,90E+00	7,00E+00
[GI tract] Colon	6,52E+00	1,20E+01
[GI tract] Defecography	8,11E+00	1,20E+01
[Biliary tract] Retrograde cholangiography	1,86E+01	1,80E+00
[Biliary tract] Operative cholangiography	9,73E-01	1,80E+00
[Biliary tract] Intravenous cholangiography	3,52E+00	1,80E+00
[Biliary tract] T drain cholangiography	2,85E+00	2,60E+00
[Biliary tract] Transhepatic cholangiography	1,80E+00	8,10E+00
[Biliary tract] ERCP	1,86E+01	5,49E+00
[Biliary tract] Retrograde pancreatography	1,86E+01	1,80E+00
[Biliary tract] Cholecystography	4,04E+00	3,10E+00
[Uro-genital] Intravenous urography	2,88E+00	2,10E+00
[Uro-genital] Retrograde pyelography	9,25E+00	2,30E+00
[Uro-genital] Nephrostography	1,60E+00	1,60E+00
[Uro-genital] Retrograde cystography	2,36E+00	1,80E+00
[Uro-genital] Micturitional cysto-urethrography	6,17E+00	3,50E+00
[Uro-genital] Urethrography	9,25E+00	1,10E+00
[Uro-genital] Hysterosalpingography	2,10E+00	2,50E+00
[Spinal cord] Cervical myelography	6,00E-01	6,00E-01
[Spinal cord] Thoracic myelography	3,88E+00	3,88E+00
[Spinal cord] Lumbar myelography	6,50E+00	6,50E+00
[Spinal cord] Sacral myelography	3,99E+00	3,99E+00
[Spinal cord] Whole spine myelography	1,10E+01	1,10E+01
[Joints] Temporal-mandibular joint arthrography	7,78E-01	7,78E-01
[Joints] Shoulder arthrography	1,30E-01	1,30E-01
[Joints] Hip arthrography	3,16E-02	3,20E-02
[Joints] Elbow arthrography	5,54E-03	6,00E-03
[Joints] Wrist arthrography	9,21E-04	1,00E-03
[Joints] Knee arthrography	5,58E-02	8,70E-02
[Joints] Ankle arthrography	2,83E-03	3,00E-03
[Other] Colonoscopy	4,84E+00	4,84E+00
[Other] Chest fluoroscopy	3,19E+00	3,19E+00
[Other] Abdominal fluoroscopy	4,03E+00	4,03E+00
[Other] Limbs/Joints fluoroscopy	2,08E-01	2,08E-01

Table 3: Effective doses for therapeutic interventions

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀
[Head & neck] Cerebral dilatation/stenting	3,84E+00	1,65E+01
[Head & neck] Cerebral embolisation (AVM, a.aneurysm, a.tumour)	3,30E+00	3,29E+01
[Head & neck] Cerebral thrombolysis	1,69E+00	1,69E+00
[Head & neck] Head & neck puncture	1,68E-01	1,70E-01
[Chest] Coronary dilatation/stenting (PTCA)	1,08E+01	1,70E+01
[Chest] Cardiac pacemaker fitting (temporary or permanent)	6,10E+00	2,80E+00
[Chest] Central venous line fitting (Hickman line)	8,89E+00	2,50E-01
[Chest] Cardiac thermo-ablation	3,16E+01	2,03E+01
[Chest] Valvuloplasty	2,42E+01	2,93E+01
[Chest] Inferior vena cava (IVC) filter fitting	8,89E+00	3,60E+00
[Chest] Oesophagus dilatation/stenting	9,69E+00	1,50E+00
[Chest] Thoracic dilatation/stenting	1,36E+01	2,94E+01
[Chest] Thoracic embolisation	4,85E+01	2,94E+01
[Chest] Thoracic thrombolysis	2,58E+01	2,94E+01
[Chest] Thoracic region biopsy	1,98E+01	4,00E-01
[Chest] Electrophysiology	3,16E+01	3,30E+00
[Abdomen] Bile duct dilatation/stenting	1,31E+01	1,40E+01
[Abdomen] Bile duct drainage	1,69E+01	3,82E+01
[Abdomen] Bile duct stone extraction	1,91E+01	7,00E+00
[Abdomen] Renal artery dilatation/stenting	1,21E+01	1,20E+01
[Abdomen] Renal drainage	1,89E+01	1,89E+01
[Abdomen] Lithotripsy	3,79E+00	3,79E+00
[Abdomen] Nephrostomy	1,35E+01	1,35E+01
[Abdomen] TIPS (liver)	6,46E+01	6,46E+01
[Abdomen] Abdominal dilatation/stenting	2,25E+01	1,50E+00
[Abdomen] Abdominal embolisation	3,68E+01	3,68E+01
[Abdomen] Abdominal thrombolysis	1,65E+01	1,65E+01
[Abdomen] Abdominal region biopsy		1,50E+00
[Pelvis] Pelvic vessel dilatation	1,45E+01	1,45E+01
[Pelvis] Pelvic vessel embolisation	3,72E+01	3,72E+01
[Pelvis] Pelvic vessel thrombolysis	1,14E+01	1,14E+01
[Limbs] Upper limb dilatation	1,78E-01	1,80E-01
[Limbs] Upper limb embolisation	4,01E-01	4,00E-01
[Limbs] Upper limb thrombolysis	1,34E-01	1,30E-01
[Limbs] Popliteal dilatation (behind knee)	2,33E+00	2,33E+00
[Limbs] Lower limb dilatation	1,49E+00	9,00E-01
[Limbs] Lower limb embolisation	4,17E+00	9,00E-01
[Limbs] Lower limb thrombolysis	2,38E+00	9,00E-01
[Limbs] Limbs biopsy	9,00E-01	9,00E-01
[Other] Thoracic puncture	1,68E-01	2,29E+00
[Other] Abdominal puncture	2,54E+00	2,54E+00
[Other] Extremities puncture	1,47E-01	1,47E-01
[Other] Vertebroplasty	1,26E+01	1,26E+01

Table 4: Effective doses for diagnostic interventions

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀
[Blood vessels] Cerebral angiography	2,70E+00	5,00E+00
[Blood vessels] Petrous phlebography	1,28E+00	1,28E+00
[Blood vessels] Coronary angiography	9,24E+00	1,12E+01
[Blood vessels] Thoracic aortography	2,33E+00	5,00E+00
[Blood vessels] Bronchial arteriography	1,10E+01	5,00E+00
[Blood vessels] Pulmonary arteriography	2,83E+01	5,00E+00
[Blood vessels] Upper venacavography	5,15E+00	5,00E+00
[Blood vessels] Abdominal aortography	6,67E+00	1,20E+01
[Blood vessels] Renal arteriography	1,27E+01	6,00E+00
[Blood vessels] Mesenteric arteriography	1,00E+01	1,00E+01
[Blood vessels] Lower venacavography	1,33E+01	1,20E+01
[Blood vessels] Renal phlebography	5,08E+00	6,00E+00
[Blood vessels] Suprarenal phlebography	5,08E+00	6,00E+00
[Blood vessels] Pelvic arteriography	1,63E+01	1,63E+01
[Blood vessels] Ovarian phlebography	3,04E+00	3,04E+00
[Blood vessels] Spermatic phlebography	6,24E+00	6,24E+00
[Blood vessels] Upper limb arteriography	3,62E+00	8,00E+00
[Blood vessels] Upper limb phlebography	4,25E-01	8,00E+00
[Blood vessels] Lower limb arteriography	6,08E+00	8,00E+00
[Blood vessels] Lower limb phlebography	2,28E+00	8,00E+00
[Lymph vessels] Thoracic lymphangiography	1,69E-01	1,70E-01
[Lymph vessels] Abdominal lymphangiography	2,67E-01	2,70E-01
[Lymph vessels] Pelvic lymphangiography	1,84E-01	1,80E-01
[Lymph vessels] Upper limb lymphangiography	1,90E-03	1,90E-03
[Lymph vessels] Lower limb lymphangiography	7,33E-02	7,00E-02

Table 5: Effective doses for CT examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀	E 2008 ICRP₁₀₃
[Head] Orbits	2,01E+00	1,24E+00	1,55E+00
[Head] Temporal bones	2,41E+00	8,00E-01	1,00E+00
[Head] Temporo-mandibular joint	1,23E+00	8,00E-01	1,00E+00
[Head] Sella turcica	1,23E+00	8,00E-01	1,00E+00
[Head] Face	1,09E+00	5,00E-01	5,00E-01
[Head] Dental	3,20E+00	5,00E-01	5,00E-01
[Head] Cerebrum	2,14E+00	2,14E+00	2,60E+00
[Head] Posterior fossa	2,14E+00	2,14E+00	2,60E+00
[Head] Brain vascular	2,14E+00	2,14E+00	2,60E+00
[Head] Pituitary gland	1,23E+00	2,14E+00	2,60E+00
[Head] Sinuses	1,09E+00	5,00E-01	5,00E-01
[Head] Internal auditory meatus	2,41E+00	5,00E-01	5,00E-01
[Head] Nasal cavity	1,23E+00	5,00E-01	5,00E-01
[Head] Mouth	3,20E+00	5,00E-01	5,00E-01
[Neck] Cervical spine	2,31E+00	2,89E+00	2,57E+00
[Neck] Neck	3,42E+00	2,78E+00	2,57E+00
[Neck] Larynx	3,42E+00	2,78E+00	2,57E+00
[Neck] Pharynx	3,42E+00	2,78E+00	2,57E+00
[Neck] Neck vascular	3,09E+00	3,09E+00	2,86E+00
[Chest + abdomen] Thoracic spine	5,21E+00	5,40E+00	5,76E+00
[Chest + abdomen] Mediastinum	6,48E+00	5,40E+00	5,76E+00
[Chest + abdomen] Lungs standard	9,02E+00	5,40E+00	5,76E+00
[Chest + abdomen] Lungs High Resolution	9,02E+00	5,40E+00	5,76E+00
[Chest + abdomen] Heart	5,93E+00	9,30E+00	9,90E+00
[Chest + abdomen] Thoracic aorta	5,93E+00	5,66E+00	5,99E+00
[Chest + abdomen] Lungs vascular	9,02E+00	5,58E+00	5,95E+00
[Abdomen] Lumbar spine	9,37E+00	9,20E+00	9,10E+00
[Abdomen] Full abdomen	1,04E+01	1,17E+01	
[Abdomen] Upper abdomen	3,99E+00	9,42E+00	9,26E+00
[Abdomen] Liver / pancreas	4,83E+00	9,96E+00	9,79E+00
[Abdomen] Kidneys / Supra-renal glands	4,83E+00	9,96E+00	9,79E+00
[Abdomen] Enteroclysis	9,96E+00	9,96E+00	9,79E+00
[Pelvis] Hip / pelvic bone	7,16E+00	7,20E+00	5,70E+00
[Pelvis] Sacrum / coccyx	7,17E+00	7,20E+00	5,70E+00
[Pelvis] Sacro-iliac joint	7,17E+00	7,20E+00	5,70E+00
[Pelvis] Pelvimetry (obstetric)	7,16E+00	7,20E+00	5,70E+00
[Pelvis] Pelvis (soft tissues/vascular)	7,16E+00	9,84E+00	7,58E+00
[Pelvis] Virtual CT colonoscopy	8,85E+00	8,85E+00	
[Neck + chest + abdomen] Full spine	1,67E+01	1,67E+01	
[Chest + abdomen] Chest & abdomen	1,08E+01	1,08E+01	1,06E+01
[Abdomen + pelvis] Abdomen & pelvis	1,17E+01	1,17E+01	1,43E+01
[Abdomen + pelvis] CT urography	6,40E+00	6,40E+00	5,00E+00
[Chest + abdomen + pelvis] Whole trunk	1,14E+01	1,14E+01	1,12E+01

[Limbs] Shoulder	2,65E+00	1,80E+00
[Limbs] Elbow	1,49E-02	1,80E+00
[Limbs] Wrist	9,11E-03	1,80E+00
[Limbs] Hand	9,11E-03	1,80E+00
[Limbs] Thigh	2,20E+00	1,00E-01
[Limbs] Knee	6,25E-02	1,00E-01
[Limbs] Lower leg	1,94E-02	1,00E-01
[Limbs] Calcaneum	1,94E-02	1,00E-01
[Limbs] Ankle	1,94E-02	1,00E-01
[Limbs] Foot	1,94E-02	1,00E-01
[Other] CT angiography	3,90E+01	3,90E+01
[Other] CT intervention	3,90E+01	3,90E+01

Table 6: Effective doses for dental examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀
[Teeth & gums] Total Periapical	5,41E-02	5,00E-03
[Teeth & gums] Total Bitewing	6,74E-03	1,00E-02
[Teeth & gums] Occlusal	7,20E-03	2,50E-02
[Teeth & gums] Panoramic radiography (OPG)	5,99E-02	2,40E-02
[Teeth & gums] CT (mandible/maxilla)	7,28E-01	5,00E-01
[Head] Skull		6,10E-02
[Head] Teleradiography	6,05E-02	6,10E-02
[Limbs] Hand	3,00E-04	3,00E-04

Table 7: Effective doses for mammography examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀
[Breast] Symptomatic (1-2 views / 1 or both breasts)	1,00E-01	1,60E-01
[Breast] Screening (1-2 views / both breasts)	2,00E-01	1,60E-01

Table 8: Effective doses for bone densitometry examinations

Type of examination	E 1998 ICRP₆₀	E 2008 ICRP₆₀	E 2008 ICRP₁₀₃
[Abdomen] Lumbar spine	1,46E-03	4,00E-03	4,00E-03
[Pelvis] Hip	3,25E-03	3,00E-03	2,00E-03
[Limbs] Upper or lower limb	9,73E-05	1,00E-04	
[Other] Whole body	3,08E-03	3,34E-03	

Appendix 5:

Geographic distribution of the
healthcare providers (total and
respondents)

Region	Cantons	ZDI			CDI			ADI			ZDI	+CDI	+ADI
		Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate
1. Lemanic region	VD / VS / GE	689	277	0.40	26	13	0.50	713	285	0.40	1428	575	0.40
2. Midland space	BE / FR / SO / NE / JU	759	320	0.42	29	19	0.66	1038	442	0.43	1826	781	0.43
3. North-West Switzerland	BS / BL / AG	449	198	0.44	14	9	0.64	621	270	0.43	1084	477	0.44
4. Zurich	ZH	622	232	0.37	21	17	0.81	823	349	0.42	1466	598	0.41
5. Eastern Switzerland	GL / SH / AR / AI / SG / GR / TG	481	208	0.43	15	13	0.87	727	330	0.45	1223	551	0.45
6. Central Switzerland	LU / UR / SZ / OW / NW / ZG	324	136	0.42	16	10	0.63	444	183	0.41	784	329	0.42
7. Tessin	TI	202	95	0.47	3	3	1.00	231	104	0.45	436	202	0.46
Switzerland		3526	1466	0.42	124	84	0.68	4597	1963	0.43	8247	3513	0.43

Region	Cantons	RADIO			UNISP			KASP			BEZSP		
		Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate
1. Lemanic region	VD / VS / GE	31	11	0.35	23	10	0.43	2	0	-	22	13	0.59
2. Midland space	BE / FR / SO / NE / JU	17	9	0.53	14	5	0.36	5	3	0.60	42	33	0.79
3. North-West Switzerland	BS / BL / AG	10	5	0.50	7	4	0.57	12	10	0.83	9	7	0.78
4. Zurich	ZH	11	8	0.73	9	7	0.78	2	1	0.50	9	6	0.67
5. Eastern Switzerland	GL / SH / AR / AI / SG / GR / TG	11	3	0.27	0	0	-	30	18	0.60	19	14	0.74
6. Central Switzerland	LU / UR / SZ / OW / NW / ZG	2	2	1.00	0	0	-	11	8	0.73	4	2	0.50
7. Tessin	TI	3	1	0.33	0	0	-	0	0	-	6	5	0.83
Switzerland		85	39	0.46	53	26	0.49	62	40	0.65	111	80	0.72

Region	Cantons	STASP			PRISP			KRANKALT			All	Hosp	
		Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate	Total	Respond	Rate
1. Lemanic region	VD / VS / GE	0	0	-	38	28	0.74	0	0	-	85	51	0.60
2. Midland space	BE / FR / SO / NE / JU	5	2	0.40	20	11	0.55	5	3	0.60	91	57	0.63
3. North-West Switzerland	BS / BL / AG	2	2	1.00	21	13	0.62	0	0	-	51	36	0.71
4. Zurich	ZH	7	6	0.86	14	6	0.43	1	1	1.00	42	27	0.64
5. Eastern Switzerland	GL / SH / AR / AI / SG / GR / TG	0	0	-	20	16	0.80	1	0	-	70	48	0.69
6. Central Switzerland	LU / UR / SZ / OW / NW / ZG	0	0	-	7	3	0.43	0	0	-	22	13	0.59
7. Tessin	TI	0	0	-	12	10	0.83	4	2	0.50	22	17	0.77
Switzerland		14	10	0.71	132	87	0.66	11	6	0.55	383	249	0.65

ZDI: Dental – CDI: Chiropractic – ADI: Medical – RADIO: Radiology institutes – UNISP: Departments at university hospitals – KASP: Departments at canton hospitals – BEZSP: District hospitals – STASP: City hospitals – PRISP: Private hospitals and clinics – KRANKALT: Homes for the elderly.

Appendix 6: Sensitivity analysis

A hypothesis on a + or - 20% difference in the frequencies of examinations that would be performed by the non-participants is considered.

	Established	80%		120%	
	Frequencies	Frequency	Ratio	Frequency	Ratio
Radiography	6.00E+06	5.59E+06	0.93	6.63E+06	1.10
Fluoroscopy	1.53E+05	1.41E+05	0.93	1.65E+05	1.08
Therapeutic intervention	4.60E+04	4.44E+04	0.97	4.84E+04	1.05
Diagnostic intervention	5.56E+04	5.29E+04	0.95	5.81E+04	1.04
Computed tomography	7.80E+05	7.40E+05	0.95	8.25E+05	1.06
Dental radiology	5.43E+06	4.80E+06	0.88	6.06E+06	1.12
Mammography	3.87E+05	3.64E+05	0.94	4.10E+05	1.06
Bone densitometry	1.17E+05	1.10E+05	0.94	1.24E+05	1.06
Total	1.30E+07	1.18E+07	0.91	1.43E+07	1.10

Appendix 7:
Examination frequency
by healthcare providers

Table 1: Frequency for radiography examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head] Orbits	0,00E+00	9,72E+03	1,40E+02	1,63E+04	1,39E+03	2,75E+04
[Head] Temporal bones	0,00E+00	6,07E+03	0,00E+00	2,07E+03	6,43E+02	8,78E+03
[Head] Sphenoid bone	0,00E+00	2,41E+03	0,00E+00	1,13E+04	5,35E+01	1,38E+04
[Head] Nose	0,00E+00	3,28E+03	0,00E+00	6,87E+03	1,70E+02	1,03E+04
[Head] Sinuses	0,00E+00	5,72E+04	0,00E+00	1,63E+04	1,90E+03	7,53E+04
[Head] Zygomas	0,00E+00	1,59E+02	0,00E+00	3,79E+02	1,49E+02	6,88E+02
[Head] Temporo-mandibular joint	0,00E+00	8,71E+02	0,00E+00	6,12E+02	1,44E+02	1,63E+03
[Head] Cervico-occipital hinge	0,00E+00	3,56E+02	9,25E+01	1,95E+03	5,59E+01	2,46E+03
[Head] Maxilla	0,00E+00	7,49E+02	0,00E+00	3,79E+02	1,65E+02	1,29E+03
[Head] Mandible	0,00E+00	5,86E+03	0,00E+00	1,50E+03	1,77E+02	7,53E+03
[Head] Cephalometry	0,00E+00	7,41E+01	0,00E+00	1,23E+02	0,00E+00	1,97E+02
[Head] Dacryocystography	0,00E+00	6,37E+02	0,00E+00	2,23E+03	7,97E+01	2,94E+03
[Head] Sialography	0,00E+00	0,00E+00	0,00E+00	1,02E+02	8,30E+00	1,11E+02
[Head] Eyes	0,00E+00	6,37E+02	0,00E+00	1,28E+01	0,00E+00	6,50E+02
[Neck] Cervical spine	0,00E+00	7,96E+04	2,50E+04	8,00E+04	1,05E+04	1,95E+05
[Neck] Larynx	0,00E+00	2,81E+03	0,00E+00	3,77E+03	3,15E+02	6,89E+03
[Neck] Pharynx	0,00E+00	2,55E+03	0,00E+00	3,50E+03	9,46E+02	6,99E+03
[Neck] Trachea	0,00E+00	2,72E+03	0,00E+00	6,85E+02	2,84E+03	6,25E+03
[Chest / Thorax] Thoracic spine	0,00E+00	2,88E+04	3,08E+03	4,69E+04	7,41E+03	8,62E+04
[Chest / Thorax] Shoulder blades / scapulae	0,00E+00	8,95E+04	4,50E+02	1,05E+05	1,16E+04	2,07E+05
[Chest / Thorax] Collar bone(s) / clavicle(s)	0,00E+00	1,53E+04	6,97E+01	2,32E+04	1,88E+03	4,05E+04
[Chest / Thorax] Acromio-clavicular joint	0,00E+00	2,93E+04	8,28E+02	7,90E+03	1,37E+03	3,94E+04

[Chest / Thorax] Sterno-clavicular joint	0,00E+00	2,27E+03	2,47E+02	3,36E+03	1,25E+02	6,00E+03
[Chest / Thorax] Manubrio-sternal joint	0,00E+00	1,24E+01	0,00E+00	6,43E+01	0,00E+00	7,67E+01
[Chest / Thorax] Sternum	0,00E+00	1,12E+03	0,00E+00	2,20E+03	1,31E+02	3,45E+03
[Chest / Thorax] Ribs	0,00E+00	2,54E+04	8,68E+01	9,16E+03	9,11E+02	3,55E+04
[Chest / Thorax] Lung	0,00E+00	6,94E+05	1,16E+02	1,02E+06	3,60E+04	1,75E+06
[Chest / Thorax] Thoracic inlet	0,00E+00	4,68E+03	2,33E+01	2,64E+03	0,00E+00	7,34E+03
[Chest / Thorax] Bronchography	0,00E+00	5,34E+02	0,00E+00	0,00E+00	0,00E+00	5,34E+02
[Abdomen] Lumbar spine	0,00E+00	1,43E+05	2,14E+04	1,49E+05	2,58E+04	3,40E+05
[Abdomen] Lumbo-sacral joint	0,00E+00	6,30E+02	1,69E+02	3,09E+03	2,72E+02	4,16E+03
[Abdomen] Abdomen (plain film, patient supine or erect)	0,00E+00	2,31E+04	9,26E+00	1,24E+05	3,78E+03	1,51E+05
[Pelvis] Ilium / ischium / pubis	0,00E+00	6,23E+04	6,93E+03	1,72E+05	1,56E+04	2,56E+05
[Pelvis] Sacrum	0,00E+00	5,66E+03	1,26E+02	8,10E+03	9,81E+02	1,49E+04
[Pelvis] Sacro-iliac joint	0,00E+00	3,68E+03	2,07E+03	4,21E+03	4,19E+03	1,41E+04
[Pelvis] Coccyx	0,00E+00	9,14E+02	7,78E+01	4,84E+03	1,11E+02	5,95E+03
[Pelvis] Pelvimetry (obstetric)	0,00E+00	0,00E+00	0,00E+00	4,45E+01	0,00E+00	4,45E+01
[Pelvis] Hips (one or both)	0,00E+00	5,29E+04	5,01E+02	1,16E+05	1,14E+04	1,81E+05
[Pelvis] Pelvis (soft tissue)	0,00E+00	1,82E+03	1,39E+03	4,67E+03	2,86E+02	8,16E+03
[Limbs] Upper arm (humerus)	0,00E+00	1,60E+04	1,17E+02	2,93E+04	1,43E+03	4,68E+04
[Limbs] Elbow	0,00E+00	4,28E+04	1,29E+02	6,51E+04	2,20E+03	1,10E+05
[Limbs] Forearm (radius & ulna)	0,00E+00	1,68E+04	3,01E+01	3,89E+04	8,23E+02	5,66E+04
[Limbs] Wrist (scaphoid)	0,00E+00	1,33E+05	1,91E+02	1,35E+05	6,32E+03	2,74E+05
[Limbs] Hand	0,00E+00	1,14E+05	2,50E+02	1,04E+05	7,00E+03	2,25E+05
[Limbs] Fingers & thumbs	0,00E+00	1,24E+05	7,62E+01	9,89E+04	5,58E+03	2,29E+05
[Limbs] Femur	0,00E+00	7,45E+03	1,41E+01	2,56E+04	1,46E+03	3,45E+04
[Limbs] Knee joint	0,00E+00	2,88E+05	6,19E+02	2,34E+05	2,24E+04	5,45E+05

[Limbs] Knee cap (patella)	0,00E+00	2,31E+04	1,17E+02	3,08E+04	1,50E+03	5,55E+04
[Limbs] Lower leg (tibia & fibula)	0,00E+00	1,72E+04	9,86E+00	4,41E+04	2,72E+03	6,41E+04
[Limbs] Ankle	0,00E+00	1,59E+05	2,37E+02	1,73E+05	8,11E+03	3,41E+05
[Limbs] Foot	0,00E+00	1,63E+05	2,76E+02	1,65E+05	1,04E+04	3,38E+05
[Limbs] Calcaneum (heel)	0,00E+00	1,79E+04	8,38E+01	1,65E+04	6,14E+03	4,06E+04
[Limbs] Toes	0,00E+00	5,85E+04	9,04E+01	2,61E+04	2,08E+03	8,68E+04
[Limbs] Whole leg (Leg length)	0,00E+00	1,04E+03	0,00E+00	1,40E+04	8,50E+02	1,59E+04
[Trunk] Scoliosis (Whole spine)	0,00E+00	2,37E+03	6,67E+01	1,12E+04	3,20E+02	1,40E+04
[Head & trunk] Whole skeleton (Skeletal survey)	0,00E+00	1,87E+03	0,00E+00	9,48E+01	0,00E+00	1,96E+03
Total	0,00E+00	2,55E+06	6,51E+04	3,17E+06	2,21E+05	6,00E+06

Table 2: Frequency for fluoroscopy examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[GI tract] Oesophagus	0,00E+00	1,27E+03	0,00E+00	1,17E+04	1,52E+03	1,45E+04
[GI tract] Stomach & duodenum	0,00E+00	3,73E+01	0,00E+00	8,40E+03	6,31E+02	9,07E+03
[GI tract] Small intestine	0,00E+00	3,73E+01	0,00E+00	2,24E+03	2,27E+02	2,50E+03
[GI tract] Enteroclysis	0,00E+00	0,00E+00	0,00E+00	4,35E+02	2,72E+02	7,07E+02
[GI tract] Colon	0,00E+00	4,78E+02	0,00E+00	4,29E+03	4,79E+02	5,25E+03
[GI tract] Defecography	0,00E+00	0,00E+00	0,00E+00	1,39E+03	6,55E+02	2,04E+03
[Biliary tract] Retrograde cholangiography	0,00E+00	0,00E+00	0,00E+00	1,72E+03	0,00E+00	1,72E+03
[Biliary tract] Operative cholangiography	0,00E+00	0,00E+00	0,00E+00	8,82E+02	0,00E+00	8,82E+02
[Biliary tract] Intravenous cholangiography	0,00E+00	0,00E+00	0,00E+00	2,72E+02	0,00E+00	2,72E+02
[Biliary tract] T drain cholangiography	0,00E+00	0,00E+00	0,00E+00	9,60E+02	0,00E+00	9,60E+02
[Biliary tract] Transhepatic cholangiography	0,00E+00	0,00E+00	0,00E+00	3,23E+02	0,00E+00	3,23E+02
[Biliary tract] ERCP	0,00E+00	4,64E+02	0,00E+00	4,87E+03	0,00E+00	5,33E+03
[Biliary tract] Retrograde pancreatography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Biliary tract] Cholecystography	0,00E+00	0,00E+00	0,00E+00	2,52E+00	0,00E+00	2,52E+00
[Uro-genital] Intravenous urography	0,00E+00	2,93E+03	0,00E+00	5,55E+03	3,37E+02	8,81E+03
[Uro-genital] Retrograde pyelography	0,00E+00	2,56E+02	0,00E+00	6,91E+03	2,57E+00	7,17E+03
[Uro-genital] Nephrostography	0,00E+00	0,00E+00	0,00E+00	1,88E+03	0,00E+00	1,88E+03
[Uro-genital] Retrograde cystography	0,00E+00	2,32E+02	0,00E+00	5,91E+03	7,99E+00	6,15E+03
[Uro-genital] Micturitional cysto-urethrography	0,00E+00	2,95E+02	0,00E+00	9,43E+03	1,08E+02	9,83E+03
[Uro-genital] Urethrography	0,00E+00	3,77E+02	0,00E+00	1,93E+03	0,00E+00	2,31E+03
[Uro-genital] Hysterosalpingography	0,00E+00	0,00E+00	0,00E+00	3,53E+03	1,22E+03	4,74E+03
[Spinal cord] Cervical myelography	0,00E+00	0,00E+00	0,00E+00	2,21E+02	3,10E+00	2,24E+02

[Spinal cord] Thoracic myelography	0,00E+00	0,00E+00	0,00E+00	2,74E+02	3,10E+00	2,77E+02
[Spinal cord] Lumbar myelography	0,00E+00	4,86E+01	0,00E+00	3,17E+03	1,89E+02	3,41E+03
[Spinal cord] Sacral myelography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Spinal cord] Whole spine myelography	0,00E+00	0,00E+00	0,00E+00	4,01E+01	1,24E+01	5,25E+01
[Joints] Temporal-mandibular joint arthrography	0,00E+00	7,29E+01	0,00E+00	0,00E+00	4,65E+00	7,75E+01
[Joints] Shoulder arthrography	0,00E+00	8,38E+02	0,00E+00	2,47E+04	1,52E+04	4,08E+04
[Joints] Hip arthrography	0,00E+00	1,13E+03	0,00E+00	3,35E+03	1,87E+03	6,36E+03
[Joints] Elbow arthrography	0,00E+00	0,00E+00	0,00E+00	4,15E+02	6,98E+01	4,84E+02
[Joints] Wrist arthrography	0,00E+00	0,00E+00	0,00E+00	1,51E+03	1,15E+03	2,66E+03
[Joints] Knee arthrography	0,00E+00	4,37E+02	0,00E+00	2,54E+03	1,23E+03	4,21E+03
[Joints] Ankle arthrography	0,00E+00	0,00E+00	0,00E+00	8,74E+02	3,49E+02	1,22E+03
[Other] Colonoscopy	0,00E+00	3,98E+02	0,00E+00	7,81E+02	0,00E+00	1,18E+03
[Other] Chest fluoroscopy	0,00E+00	0,00E+00	0,00E+00	2,69E+03	2,39E+01	2,71E+03
[Other] Abdominal fluoroscopy	0,00E+00	0,00E+00	0,00E+00	1,44E+03	0,00E+00	1,44E+03
[Other] Limbs/Joints fluoroscopy	0,00E+00	0,00E+00	0,00E+00	3,02E+03	6,46E+01	3,09E+03
Total	0,00E+00	9,32E+03	0,00E+00	1,18E+05	2,57E+04	1,53E+05

Table 3: Frequency for therapeutic interventions according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head & neck] Cerebral dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,10E+02	0,00E+00	3,10E+02
[Head & neck] Cerebral embolisation (AVM, a.aneurysm, a.tumour)	0,00E+00	0,00E+00	0,00E+00	4,38E+02	0,00E+00	4,38E+02
[Head & neck] Cerebral thrombolysis	0,00E+00	0,00E+00	0,00E+00	2,75E+02	0,00E+00	2,75E+02
[Head & neck] Head & neck puncture	0,00E+00	0,00E+00	0,00E+00	4,48E+02	0,00E+00	4,48E+02
[Chest] Coronary dilatation/stenting (PTCA)	0,00E+00	0,00E+00	0,00E+00	1,82E+04	0,00E+00	1,82E+04
[Chest] Cardiac pacemaker fitting (temporary or permanent)	0,00E+00	0,00E+00	0,00E+00	2,60E+03	0,00E+00	2,60E+03
[Chest] Central venous line fitting (Hickman line)	0,00E+00	0,00E+00	0,00E+00	3,93E+02	0,00E+00	3,93E+02
[Chest] Cardiac thermo-ablation	0,00E+00	0,00E+00	0,00E+00	7,72E+02	0,00E+00	7,72E+02
[Chest] Valvuloplasty	0,00E+00	0,00E+00	0,00E+00	1,37E+02	0,00E+00	1,37E+02
[Chest] Inferior vena cava (IVC) filter fitting	0,00E+00	0,00E+00	0,00E+00	2,64E+02	0,00E+00	2,64E+02
[Chest] Oesophagus dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,43E+02	0,00E+00	3,43E+02
[Chest] Thoracic dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,89E+02	0,00E+00	3,89E+02
[Chest] Thoracic embolisation	0,00E+00	0,00E+00	0,00E+00	1,56E+01	0,00E+00	1,56E+01
[Chest] Thoracic thrombolysis	0,00E+00	0,00E+00	0,00E+00	2,21E+01	0,00E+00	2,21E+01
[Chest] Thoracic region biopsy	0,00E+00	0,00E+00	0,00E+00	6,16E+01	0,00E+00	6,16E+01
[Chest] Electrophysiology	0,00E+00	0,00E+00	0,00E+00	1,10E+03	0,00E+00	1,10E+03
[Abdomen] Bile duct dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	6,01E+02	0,00E+00	6,01E+02
[Abdomen] Bile duct drainage	0,00E+00	0,00E+00	0,00E+00	1,06E+03	0,00E+00	1,06E+03
[Abdomen] Bile duct stone extraction	0,00E+00	0,00E+00	0,00E+00	2,05E+02	0,00E+00	2,05E+02
[Abdomen] Renal artery dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,17E+02	0,00E+00	3,17E+02
[Abdomen] Renal drainage	0,00E+00	0,00E+00	0,00E+00	5,48E+02	0,00E+00	5,48E+02

[Abdomen] Lithotripsy	0,00E+00	0,00E+00	0,00E+00	2,95E+02	0,00E+00	2,95E+02
[Abdomen] Nephrostomy	0,00E+00	0,00E+00	0,00E+00	1,08E+02	1,35E+01	1,09E+02
[Abdomen] TIPS (liver)	0,00E+00	0,00E+00	0,00E+00	1,68E+02	0,00E+00	1,68E+02
[Abdomen] Abdominal dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	2,12E+03	0,00E+00	2,12E+03
[Abdomen] Abdominal embolisation	0,00E+00	0,00E+00	0,00E+00	6,31E+02	0,00E+00	6,31E+02
[Abdomen] Abdominal thrombolysis	0,00E+00	0,00E+00	0,00E+00	1,37E+01	0,00E+00	1,37E+01
[Abdomen] Abdominal region biopsy	0,00E+00	0,00E+00	0,00E+00	1,70E+02	0,00E+00	1,70E+02
[Pelvis] Pelvic vessel dilatation	0,00E+00	0,00E+00	0,00E+00	1,21E+03	0,00E+00	1,21E+03
[Pelvis] Pelvic vessel embolisation	0,00E+00	0,00E+00	0,00E+00	4,49E+02	0,00E+00	4,49E+02
[Pelvis] Pelvic vessel thrombolysis	0,00E+00	0,00E+00	0,00E+00	1,55E+01	0,00E+00	1,55E+01
[Limbs] Upper limb dilatation	0,00E+00	0,00E+00	0,00E+00	6,98E+02	0,00E+00	6,98E+02
[Limbs] Upper limb embolisation	0,00E+00	0,00E+00	0,00E+00	1,20E+02	0,00E+00	1,20E+02
[Limbs] Upper limb thrombolysis	0,00E+00	0,00E+00	0,00E+00	1,69E+01	0,00E+00	1,69E+01
[Limbs] Popliteal dilatation (behind knee)	0,00E+00	0,00E+00	0,00E+00	4,80E+02	0,00E+00	4,80E+02
[Limbs] Lower limb dilatation	0,00E+00	0,00E+00	0,00E+00	6,44E+03	0,00E+00	6,44E+03
[Limbs] Lower limb embolisation	0,00E+00	0,00E+00	0,00E+00	3,55E+01	0,00E+00	3,55E+01
[Limbs] Lower limb thrombolysis	0,00E+00	0,00E+00	0,00E+00	4,07E+02	0,00E+00	4,07E+02
[Limbs] Limbs biopsy	0,00E+00	0,00E+00	0,00E+00	0,00E+00	<u>9,00E-01</u>	1,00E+00
[Other] Thoracic puncture	0,00E+00	0,00E+00	0,00E+00	6,54E+02	<u>0,00E+00</u>	6,54E+02
[Other] Abdominal puncture	0,00E+00	0,00E+00	0,00E+00	1,20E+03	<u>0,00E+00</u>	1,20E+03
[Other] Extremities puncture	0,00E+00	0,00E+00	0,00E+00	8,94E+02	<u>0,00E+00</u>	8,94E+02
[Other] Vertebroplasty	0,00E+00	0,00E+00	0,00E+00	1,26E+02	0,00E+00	1,26E+02
Total	0,00E+00	0,00E+00	0,00E+00	4,48E+04	1,44E+01	4,60E+04

Table 4: Frequency for diagnostic interventions according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Blood vessels] Cerebral angiography	0,00E+00	0,00E+00	0,00E+00	3,32E+03	0,00E+00	3,32E+03
[Blood vessels] Petrous phlebography	0,00E+00	0,00E+00	0,00E+00	3,76E+01	0,00E+00	3,76E+01
[Blood vessels] Coronary angiography	0,00E+00	0,00E+00	0,00E+00	3,42E+04	4,00E+00	3,42E+04
[Blood vessels] Thoracic aortography	0,00E+00	0,00E+00	0,00E+00	1,43E+03	0,00E+00	1,43E+03
[Blood vessels] Bronchial arteriography	0,00E+00	0,00E+00	0,00E+00	7,03E+02	0,00E+00	7,03E+02
[Blood vessels] Pulmonary arteriography	0,00E+00	0,00E+00	0,00E+00	1,11E+02	0,00E+00	1,11E+02
[Blood vessels] Upper venacavography	0,00E+00	0,00E+00	0,00E+00	1,11E+03	0,00E+00	1,11E+03
[Blood vessels] Abdominal aortography	0,00E+00	0,00E+00	0,00E+00	2,28E+03	0,00E+00	2,28E+03
[Blood vessels] Renal arteriography	0,00E+00	0,00E+00	0,00E+00	1,53E+03	0,00E+00	1,53E+03
[Blood vessels] Mesenteric arteriography	0,00E+00	0,00E+00	0,00E+00	3,19E+02	0,00E+00	3,19E+02
[Blood vessels] Lower venacavography	0,00E+00	0,00E+00	0,00E+00	3,19E+02	0,00E+00	3,19E+02
[Blood vessels] Renal phlebography	0,00E+00	0,00E+00	0,00E+00	2,75E+01	0,00E+00	2,75E+01
[Blood vessels] Suprarenal phlebography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Blood vessels] Pelvic arteriography	0,00E+00	0,00E+00	0,00E+00	1,44E+03	0,00E+00	1,44E+03
[Blood vessels] Ovarian phlebography	0,00E+00	0,00E+00	0,00E+00	2,32E+01	0,00E+00	2,32E+01
[Blood vessels] Spermatic phlebography	0,00E+00	0,00E+00	0,00E+00	2,50E+01	0,00E+00	2,50E+01
[Blood vessels] Upper limb arteriography	0,00E+00	0,00E+00	0,00E+00	9,56E+02	0,00E+00	9,56E+02
[Blood vessels] Upper limb phlebography	0,00E+00	0,00E+00	0,00E+00	7,24E+02	2,00E+00	7,26E+02
[Blood vessels] Lower limb arteriography	0,00E+00	0,00E+00	0,00E+00	6,38E+03	0,00E+00	6,38E+03
[Blood vessels] Lower limb phlebography	0,00E+00	0,00E+00	0,00E+00	6,33E+02	2,40E+01	6,57E+02
[Lymph vessels] Thoracic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Abdominal lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00

[Lymph vessels] Pelvic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Upper limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Lower limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Total	0,00E+00	0,00E+00	0,00E+00	5,55E+04	3,00E+01	5,56E+04

Table 5: Frequency for CT examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head] Orbits	0,00E+00	0,00E+00	0,00E+00	1,43E+04	1,18E+03	1,55E+04
[Head] Temporal bones	0,00E+00	0,00E+00	0,00E+00	2,74E+04	7,08E+02	2,81E+04
[Head] Temporo-mandibular joint	0,00E+00	0,00E+00	0,00E+00	3,63E+03	4,90E+02	4,12E+03
[Head] Sella turcica	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Face	0,00E+00	0,00E+00	0,00E+00	7,52E+03	2,18E+03	9,69E+03
[Head] Dental	1,15E+04	0,00E+00	0,00E+00	1,10E+03	2,73E+03	1,53E+04
[Head] Cerebrum	0,00E+00	0,00E+00	0,00E+00	7,51E+04	7,83E+03	8,29E+04
[Head] Posterior fossa	0,00E+00	0,00E+00	0,00E+00	2,74E+03	5,75E+01	2,80E+03
[Head] Brain vascular	0,00E+00	0,00E+00	0,00E+00	2,15E+03	5,02E+02	2,66E+03
[Head] Pituitary gland	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Sinuses	0,00E+00	0,00E+00	0,00E+00	8,20E+03	7,03E+03	1,52E+04
[Head] Internal auditory meatus	0,00E+00	0,00E+00	0,00E+00	1,47E+03	1,06E+01	1,48E+03
[Head] Nasal cavity	0,00E+00	0,00E+00	0,00E+00	5,51E+03	8,99E+02	6,41E+03
[Head] Mouth	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Neck] Cervical spine	0,00E+00	0,00E+00	0,00E+00	1,72E+04	2,12E+03	1,94E+04
[Neck] Neck	0,00E+00	0,00E+00	0,00E+00	1,53E+04	2,89E+03	1,82E+04
[Neck] Larynx	0,00E+00	0,00E+00	0,00E+00	3,04E+02	1,95E+02	5,00E+02
[Neck] Pharynx	0,00E+00	0,00E+00	0,00E+00	3,84E+01	0,00E+00	3,84E+01
[Neck] Neck vascular	0,00E+00	0,00E+00	0,00E+00	5,48E+03	2,50E+02	5,73E+03
[Chest + abdomen] Thoracic spine	0,00E+00	0,00E+00	0,00E+00	6,77E+03	6,62E+02	7,44E+03
[Chest + abdomen] Mediastinum	0,00E+00	0,00E+00	0,00E+00	8,35E+03	1,76E+01	8,37E+03
[Chest + abdomen] Lungs standard	0,00E+00	0,00E+00	0,00E+00	8,44E+04	1,66E+04	1,01E+05

[Chest + abdomen] Lungs High Resolution	0,00E+00	0,00E+00	0,00E+00	2,13E+03	1,79E+03	3,93E+03
[Chest + abdomen] Heart	0,00E+00	0,00E+00	0,00E+00	4,19E+03	7,00E+02	4,89E+03
[Chest + abdomen] Thoracic aorta	0,00E+00	0,00E+00	0,00E+00	5,14E+03	6,69E+02	5,81E+03
[Chest + abdomen] Lungs vascular	0,00E+00	0,00E+00	0,00E+00	8,20E+03	9,82E+02	9,18E+03
[Abdomen] Lumbar spine	0,00E+00	0,00E+00	0,00E+00	2,30E+04	7,34E+03	3,04E+04
[Abdomen] Full abdomen	0,00E+00	0,00E+00	0,00E+00	1,32E+05	1,93E+04	1,52E+05
[Abdomen] Upper abdomen	0,00E+00	0,00E+00	0,00E+00	1,44E+04	1,15E+04	2,59E+04
[Abdomen] Liver / pancreas	0,00E+00	0,00E+00	0,00E+00	3,43E+03	1,23E+02	3,55E+03
[Abdomen] Kidneys / Supra-renal glands	0,00E+00	0,00E+00	0,00E+00	3,67E+03	2,29E+02	3,90E+03
[Abdomen] Enteroclysis	0,00E+00	0,00E+00	0,00E+00	3,57E+01	1,36E+01	4,94E+01
[Pelvis] Hip / pelvic bone	0,00E+00	0,00E+00	0,00E+00	5,38E+03	3,94E+03	9,35E+03
[Pelvis] Sacrum / coccyx	0,00E+00	0,00E+00	0,00E+00	2,33E+03	2,13E+02	2,54E+03
[Pelvis] Sacro-iliac joint	0,00E+00	0,00E+00	0,00E+00	1,29E+03	3,28E+03	4,57E+03
[Pelvis] Pelvimetry (obstetric)	0,00E+00	0,00E+00	0,00E+00	2,32E+02	2,27E+00	2,34E+02
[Pelvis] Pelvis (soft tissues/vascular)	0,00E+00	0,00E+00	0,00E+00	5,53E+03	1,07E+03	6,60E+03
[Pelvis] Virtual CT colonoscopy	0,00E+00	0,00E+00	0,00E+00	4,28E+02	6,52E+02	1,08E+03
[Neck + chest + abdomen] Full spine	0,00E+00	0,00E+00	0,00E+00	2,27E+04	5,47E+02	2,32E+04
[Chest + abdomen] Chest & abdomen	0,00E+00	0,00E+00	0,00E+00	2,58E+04	1,02E+03	2,68E+04
[Abdomen + pelvis] Abdomen & pelvis	0,00E+00	0,00E+00	0,00E+00	3,68E+04	7,95E+02	3,76E+04
[Abdomen + pelvis] CT urography	0,00E+00	0,00E+00	0,00E+00	2,16E+03	3,88E+02	2,54E+03
[Chest + abdomen + pelvis] Whole trunk	0,00E+00	0,00E+00	0,00E+00	5,66E+03	0,00E+00	5,66E+03
[Limbs] Shoulder	0,00E+00	0,00E+00	0,00E+00	5,91E+03	1,86E+03	7,79E+03
[Limbs] Elbow	0,00E+00	0,00E+00	0,00E+00	2,14E+03	4,02E+02	2,57E+03
[Limbs] Wrist	0,00E+00	0,00E+00	0,00E+00	5,30E+03	1,10E+03	6,42E+03
[Limbs] Hand	0,00E+00	0,00E+00	0,00E+00	1,50E+03	2,22E+02	1,75E+03

[Limbs] Thigh	0,00E+00	0,00E+00	0,00E+00	3,54E+03	2,82E+02	3,82E+03
[Limbs] Knee	0,00E+00	0,00E+00	0,00E+00	7,01E+03	2,09E+03	9,10E+03
[Limbs] Lower leg	0,00E+00	0,00E+00	0,00E+00	1,39E+03	5,02E+02	1,89E+03
[Limbs] Calcaneum	0,00E+00	0,00E+00	0,00E+00	6,78E+02	2,00E+02	8,84E+02
[Limbs] Ankle	0,00E+00	0,00E+00	0,00E+00	5,37E+03	8,95E+02	6,34E+03
[Limbs] Foot	0,00E+00	0,00E+00	0,00E+00	3,33E+03	1,16E+03	4,55E+03
[Other] CT angiography	0,00E+00	0,00E+00	0,00E+00	1,20E+04	7,17E+03	1,92E+04
[Other] CT intervention	0,00E+00	0,00E+00	0,00E+00	9,72E+03	1,43E+03	1,11E+04
Total	1,15E+04	0,00E+00	0,00E+00	6,50E+05	1,18E+05	7,80E+05

Table 6: Frequency for dental examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Teeth & gums] Total Periapical	2,47E+06	1,80E+03	0,00E+00	3,47E+04	0,00E+00	2,51E+06
[Teeth & gums] Total Bitewing	2,29E+06	3,58E+02	0,00E+00	1,90E+04	0,00E+00	2,31E+06
[Teeth & gums] Occlusal	4,03E+04	8,16E+01	0,00E+00	2,19E+02	2,47E+01	4,06E+04
[Teeth & gums] Panoramic radiography (OPG)	3,50E+05	7,57E+03	0,00E+00	5,46E+04	5,51E+03	4,17E+05
[Teeth & gums] CT (mandible/maxilla)	1,22E+04	2,09E+03	0,00E+00	9,29E+02	6,66E+02	1,59E+04
[Head] Skull	3,20E+04	3,54E+02	0,00E+00	4,90E+03	1,10E+02	3,74E+04
[Head] Teleradiography	8,09E+04	8,94E+02	0,00E+00	1,19E+04	7,13E+02	9,45E+04
[Limbs] Hand	4,94E+03	4,80E+00	0,00E+00	3,55E+03	6,79E+02	9,18E+03
Total	5,28E+06	1,31E+04	0,00E+00	1,30E+05	7,70E+03	5,43E+06

Table 7: Frequency for mammography examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Breast] Symptomatic (1-2 views / 1 or both breasts)	0,00E+00	2,41E+04	0,00E+00	1,65E+05	9,45E+04	2,84E+05
[Breast] Screening (1-2 views / both breasts)	0,00E+00	5,70E+03	0,00E+00	7,24E+04	2,44E+04	1,02E+05
Total	0,00E+00	2,98E+04	0,00E+00	2,38E+05	1,19E+05	3,87E+05

Table 8: Frequency for bone densitometry examinations according to different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Abdomen] Lumbar spine	0,00E+00	1,05E+03	0,00E+00	3,39E+04	8,89E+03	4,38E+04
[Pelvis] Hip	0,00E+00	1,04E+03	0,00E+00	3,36E+04	8,88E+03	4,35E+04
[Limbs] Upper or lower limb	0,00E+00	6,33E+02	0,00E+00	2,05E+04	7,63E+03	2,87E+04
[Other] Whole body	0,00E+00	0,00E+00	0,00E+00	1,13E+03	0,00E+00	1,13E+03
Total	0,00E+00	2,72E+03	0,00E+00	8,91E+04	2,54E+04	1,17E+05

Appendix 8:
Annual collective dose
by healthcare providers

Table 1: Annual collective dose (in mSv) delivered through radiography examinations by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head] Orbits	0,00E+00	2,92E+02	4,19E+00	4,88E+02	4,18E+01	8,25E+02
[Head] Temporal bones	0,00E+00	3,64E+02	0,00E+00	1,24E+02	3,86E+01	5,27E+02
[Head] Sphenoid bone	0,00E+00	7,24E+01	0,00E+00	3,39E+02	1,61E+00	4,13E+02
[Head] Nose	0,00E+00	1,31E+02	0,00E+00	2,75E+02	6,80E+00	4,13E+02
[Head] Sinuses	0,00E+00	2,63E+03	0,00E+00	7,49E+02	8,76E+01	3,47E+03
[Head] Zygomas	0,00E+00	6,38E+00	0,00E+00	1,52E+01	5,96E+00	2,75E+01
[Head] Temporo-mandibular joint	0,00E+00	3,48E+01	0,00E+00	2,45E+01	5,77E+00	6,51E+01
[Head] Cervico-occipital hinge	0,00E+00	1,49E+01	3,89E+00	8,20E+01	2,35E+00	1,03E+02
[Head] Maxilla	0,00E+00	3,00E+01	0,00E+00	1,52E+01	6,61E+00	5,17E+01
[Head] Mandible	0,00E+00	2,34E+02	0,00E+00	6,00E+01	7,08E+00	3,01E+02
[Head] Cephalometry	0,00E+00	2,97E+00	0,00E+00	4,92E+00	0,00E+00	7,88E+00
[Head] Dacryocystography	0,00E+00	5,09E+01	0,00E+00	1,78E+02	6,37E+00	2,36E+02
[Head] Sialography	0,00E+00	0,00E+00	0,00E+00	4,10E+00	3,32E-01	4,43E+00
[Head] Eyes	0,00E+00	2,55E+01	0,00E+00	5,12E-01	0,00E+00	2,60E+01
[Neck] Cervical spine	0,00E+00	8,12E+03	2,55E+03	8,16E+03	1,07E+03	1,99E+04
[Neck] Larynx	0,00E+00	1,40E+02	0,00E+00	1,88E+02	1,58E+01	3,45E+02
[Neck] Pharynx	0,00E+00	1,27E+02	0,00E+00	1,75E+02	4,73E+01	3,49E+02
[Neck] Trachea	0,00E+00	2,72E+02	0,00E+00	6,85E+01	2,84E+02	6,25E+02
[Chest / Thorax] Thoracic spine	0,00E+00	1,25E+04	1,34E+03	2,04E+04	3,22E+03	3,75E+04
[Chest / Thorax] Shoulder blades / scapulae	0,00E+00	2,60E+03	1,30E+01	3,05E+03	3,36E+02	6,00E+03
[Chest / Thorax] Collar bone(s) / clavicle(s)	0,00E+00	4,44E+02	2,02E+00	6,73E+02	5,45E+01	1,17E+03
[Chest / Thorax] Acromio-clavicular joint	0,00E+00	8,48E+02	2,40E+01	2,29E+02	3,98E+01	1,14E+03

[Chest / Thorax] Sterno-clavicular joint	0,00E+00	6,57E+01	7,16E+00	9,74E+01	3,63E+00	1,74E+02
[Chest / Thorax] Manubrio-sternal joint	0,00E+00	3,59E-01	0,00E+00	1,87E+00	0,00E+00	2,22E+00
[Chest / Thorax] Sternum	0,00E+00	3,25E+01	0,00E+00	6,38E+01	3,79E+00	1,00E+02
[Chest / Thorax] Ribs	0,00E+00	1,27E+03	4,34E+00	4,58E+02	4,55E+01	1,78E+03
[Chest / Thorax] Lung	0,00E+00	3,50E+04	5,87E+00	5,15E+04	1,82E+03	8,84E+04
[Chest / Thorax] Thoracic inlet	0,00E+00	1,78E+02	8,84E-01	1,00E+02	0,00E+00	2,79E+02
[Chest / Thorax] Bronchography	0,00E+00	2,03E+01	0,00E+00	0,00E+00	0,00E+00	2,03E+01
[Abdomen] Lumbar spine	0,00E+00	2,43E+05	3,62E+04	2,53E+05	4,37E+04	5,76E+05
[Abdomen] Lumbo-sacral joint	0,00E+00	5,18E+02	1,39E+02	2,54E+03	2,24E+02	3,42E+03
[Abdomen] Abdomen (plain film, patient supine or erect)	0,00E+00	1,81E+04	7,22E+00	9,70E+04	2,95E+03	1,18E+05
[Pelvis] Ilium / ischium / pubis	0,00E+00	5,05E+04	5,62E+03	1,39E+05	1,26E+04	2,08E+05
[Pelvis] Sacrum	0,00E+00	6,12E+03	1,36E+02	8,75E+03	1,06E+03	1,61E+04
[Pelvis] Sacro-iliac joint	0,00E+00	3,97E+03	2,23E+03	4,55E+03	4,52E+03	1,53E+04
[Pelvis] Coccyx	0,00E+00	9,87E+02	8,40E+01	5,23E+03	1,19E+02	6,42E+03
[Pelvis] Pelvimetry (obstetric)	0,00E+00	0,00E+00	0,00E+00	2,40E+01	0,00E+00	2,40E+01
[Pelvis] Hips (one or both)	0,00E+00	5,97E+04	5,65E+02	1,31E+05	1,29E+04	2,05E+05
[Pelvis] Pelvis (soft tissue)	0,00E+00	9,82E+02	7,48E+02	2,52E+03	1,54E+02	4,41E+03
[Limbs] Upper arm (humerus)	0,00E+00	9,73E+01	7,14E-01	1,79E+02	8,70E+00	2,86E+02
[Limbs] Elbow	0,00E+00	2,57E+01	7,76E-02	3,91E+01	1,32E+00	6,62E+01
[Limbs] Forearm (radius & ulna)	0,00E+00	1,85E+01	3,31E-02	4,28E+01	9,05E-01	6,22E+01
[Limbs] Wrist (scaphoid)	0,00E+00	2,66E+01	3,83E-02	2,70E+01	1,26E+00	5,49E+01
[Limbs] Hand	0,00E+00	2,28E+01	5,00E-02	2,08E+01	1,40E+00	4,50E+01
[Limbs] Fingers & thumbs	0,00E+00	1,24E+01	7,62E-03	9,89E+00	5,58E-01	2,29E+01
[Limbs] Femur	0,00E+00	2,98E+02	5,64E-01	1,02E+03	5,83E+01	1,38E+03
[Limbs] Knee joint	0,00E+00	2,30E+02	4,95E-01	1,87E+02	1,79E+01	4,36E+02

[Limbs] Knee cap (patella)	0,00E+00	9,22E+00	4,68E-02	1,23E+01	6,01E-01	2,22E+01
[Limbs] Lower leg (tibia & fibula)	0,00E+00	6,89E+00	3,95E-03	1,76E+01	1,09E+00	2,56E+01
[Limbs] Ankle	0,00E+00	6,37E+01	9,47E-02	6,92E+01	3,25E+00	1,36E+02
[Limbs] Foot	0,00E+00	6,50E+01	1,10E-01	6,60E+01	4,16E+00	1,35E+02
[Limbs] Calcaneum (heel)	0,00E+00	7,15E+00	3,35E-02	6,60E+00	2,45E+00	1,62E+01
[Limbs] Toes	0,00E+00	2,34E+01	3,62E-02	1,04E+01	8,33E-01	3,47E+01
[Limbs] Whole leg (Leg length)	0,00E+00	2,09E+01	0,00E+00	2,81E+02	1,70E+01	3,19E+02
[Trunk] Scoliosis (Whole spine)	0,00E+00	8,05E+02	2,27E+01	3,82E+03	1,09E+02	4,75E+03
[Head & trunk] Whole skeleton (Skeletal survey)	0,00E+00	6,43E+03	0,00E+00	3,26E+02	0,00E+00	6,75E+03
Total	0,00E+00	4,58E+05	4,97E+04	7,37E+05	8,56E+04	1,33E+06

Table 2: Annual collective dose (in mSv) delivered through fluoroscopy examinations by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[GI tract] Oesophagus	0,00E+00	1,40E+03	0,00E+00	1,28E+04	1,68E+03	1,59E+04
[GI tract] Stomach & duodenum	0,00E+00	4,48E+02	0,00E+00	1,01E+05	7,57E+03	1,09E+05
[GI tract] Small intestine	0,00E+00	1,42E+02	0,00E+00	8,51E+03	8,64E+02	9,52E+03
[GI tract] Enteroclysis	0,00E+00	0,00E+00	0,00E+00	3,04E+03	1,90E+03	4,95E+03
[GI tract] Colon	0,00E+00	5,73E+03	0,00E+00	5,15E+04	5,75E+03	6,30E+04
[GI tract] Defecography	0,00E+00	0,00E+00	0,00E+00	1,67E+04	7,86E+03	2,45E+04
[Biliary tract] Retrograde cholangiography	0,00E+00	0,00E+00	0,00E+00	3,10E+03	0,00E+00	3,10E+03
[Biliary tract] Operative cholangiography	0,00E+00	0,00E+00	0,00E+00	1,59E+03	0,00E+00	1,59E+03
[Biliary tract] Intravenous cholangiography	0,00E+00	0,00E+00	0,00E+00	4,90E+02	0,00E+00	4,90E+02
[Biliary tract] T drain cholangiography	0,00E+00	0,00E+00	0,00E+00	2,50E+03	0,00E+00	2,50E+03
[Biliary tract] Transhepatic cholangiography	0,00E+00	0,00E+00	0,00E+00	2,61E+03	0,00E+00	2,61E+03
[Biliary tract] ERCP	0,00E+00	2,55E+03	0,00E+00	2,67E+04	0,00E+00	2,93E+04
[Biliary tract] Retrograde pancreatography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Biliary tract] Cholecystography	0,00E+00	0,00E+00	0,00E+00	7,81E+00	0,00E+00	7,81E+00
[Uro-genital] Intravenous urography	0,00E+00	6,15E+03	0,00E+00	1,17E+04	7,07E+02	1,85E+04
[Uro-genital] Retrograde pyelography	0,00E+00	5,88E+02	0,00E+00	1,59E+04	5,91E+00	1,65E+04
[Uro-genital] Nephrostography	0,00E+00	0,00E+00	0,00E+00	3,01E+03	0,00E+00	3,01E+03
[Uro-genital] Retrograde cystography	0,00E+00	4,18E+02	0,00E+00	1,06E+04	1,44E+01	1,11E+04
[Uro-genital] Micturitional cysto-urethrography	0,00E+00	1,03E+03	0,00E+00	3,30E+04	3,77E+02	3,44E+04
[Uro-genital] Urethrography	0,00E+00	4,14E+02	0,00E+00	2,13E+03	0,00E+00	2,54E+03
[Uro-genital] Hysterosalpingography	0,00E+00	0,00E+00	0,00E+00	8,82E+03	3,04E+03	1,19E+04
[Spinal cord] Cervical myelography	0,00E+00	0,00E+00	0,00E+00	1,32E+02	1,86E+00	1,34E+02

[Spinal cord] Thoracic myelography	0,00E+00	0,00E+00	0,00E+00	1,06E+03	1,20E+01	1,08E+03
[Spinal cord] Lumbar myelography	0,00E+00	3,16E+02	0,00E+00	2,06E+04	1,23E+03	2,21E+04
[Spinal cord] Sacral myelography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Spinal cord] Whole spine myelography	0,00E+00	0,00E+00	0,00E+00	4,41E+02	1,36E+02	5,77E+02
[Joints] Temporal-mandibular joint arthrography	0,00E+00	5,67E+01	0,00E+00	0,00E+00	3,62E+00	6,03E+01
[Joints] Shoulder arthrography	0,00E+00	1,09E+02	0,00E+00	3,22E+03	1,98E+03	5,31E+03
[Joints] Hip arthrography	0,00E+00	3,62E+01	0,00E+00	1,07E+02	5,99E+01	2,03E+02
[Joints] Elbow arthrography	0,00E+00	0,00E+00	0,00E+00	2,49E+00	4,19E-01	2,91E+00
[Joints] Wrist arthrography	0,00E+00	0,00E+00	0,00E+00	1,51E+00	1,15E+00	2,66E+00
[Joints] Knee arthrography	0,00E+00	3,80E+01	0,00E+00	2,21E+02	1,07E+02	3,67E+02
[Joints] Ankle arthrography	0,00E+00	0,00E+00	0,00E+00	2,62E+00	1,05E+00	3,67E+00
[Other] Colonoscopy	0,00E+00	1,93E+03	0,00E+00	3,78E+03	0,00E+00	5,71E+03
[Other] Chest fluoroscopy	0,00E+00	0,00E+00	0,00E+00	8,57E+03	7,64E+01	8,65E+03
[Other] Abdominal fluoroscopy	0,00E+00	0,00E+00	0,00E+00	5,80E+03	0,00E+00	5,80E+03
[Other] Limbs/Joints fluoroscopy	0,00E+00	0,00E+00	0,00E+00	6,29E+02	1,34E+01	6,42E+02
Total	0,00E+00	2,14E+04	0,00E+00	3,60E+05	3,34E+04	4,15E+05

Table 3: Annual collective dose (in mSv) delivered through therapeutic interventions by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head & neck] Cerebral dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	5,11E+03	0,00E+00	5,11E+03
[Head & neck] Cerebral embolisation (AVM, a.aneurysm, a.tumour)	0,00E+00	0,00E+00	0,00E+00	1,44E+04	0,00E+00	1,44E+04
[Head & neck] Cerebral thrombolysis	0,00E+00	0,00E+00	0,00E+00	4,65E+02	0,00E+00	4,65E+02
[Head & neck] Head & neck puncture	0,00E+00	0,00E+00	0,00E+00	7,61E+01	0,00E+00	7,61E+01
[Chest] Coronary dilatation/stenting (PTCA)	0,00E+00	0,00E+00	0,00E+00	3,10E+05	0,00E+00	3,10E+05
[Chest] Cardiac pacemaker fitting (temporary or permanent)	0,00E+00	0,00E+00	0,00E+00	7,29E+03	0,00E+00	7,29E+03
[Chest] Central venous line fitting (Hickman line)	0,00E+00	0,00E+00	0,00E+00	9,83E+01	0,00E+00	9,83E+01
[Chest] Cardiac thermo-ablation	0,00E+00	0,00E+00	0,00E+00	1,57E+04	0,00E+00	1,57E+04
[Chest] Valvuloplasty	0,00E+00	0,00E+00	0,00E+00	4,03E+03	0,00E+00	4,03E+03
[Chest] Inferior vena cava (IVC) filter fitting	0,00E+00	0,00E+00	0,00E+00	9,50E+02	0,00E+00	9,50E+02
[Chest] Oesophagus dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	5,15E+02	0,00E+00	5,15E+02
[Chest] Thoracic dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	1,14E+04	0,00E+00	1,14E+04
[Chest] Thoracic embolisation	0,00E+00	0,00E+00	0,00E+00	4,58E+02	0,00E+00	4,58E+02
[Chest] Thoracic thrombolysis	0,00E+00	0,00E+00	0,00E+00	6,50E+02	0,00E+00	6,50E+02
[Chest] Thoracic region biopsy	0,00E+00	0,00E+00	0,00E+00	2,46E+01	0,00E+00	2,46E+01
[Chest] Electrophysiology	0,00E+00	0,00E+00	0,00E+00	3,64E+03	0,00E+00	3,64E+03
[Abdomen] Bile duct dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	8,41E+03	0,00E+00	8,41E+03
[Abdomen] Bile duct drainage	0,00E+00	0,00E+00	0,00E+00	4,04E+04	0,00E+00	4,04E+04
[Abdomen] Bile duct stone extraction	0,00E+00	0,00E+00	0,00E+00	1,43E+03	0,00E+00	1,43E+03
[Abdomen] Renal artery dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,81E+03	0,00E+00	3,81E+03
[Abdomen] Renal drainage	0,00E+00	0,00E+00	0,00E+00	1,03E+04	0,00E+00	1,03E+04

[Abdomen] Lithotripsy	0,00E+00	0,00E+00	0,00E+00	1,12E+03	0,00E+00	1,12E+03
[Abdomen] Nephrostomy	0,00E+00	1,35E+01	0,00E+00	1,46E+03	0,00E+00	1,48E+03
[Abdomen] TIPS (liver)	0,00E+00	0,00E+00	0,00E+00	1,08E+04	0,00E+00	1,08E+04
[Abdomen] Abdominal dilatation/stenting	0,00E+00	0,00E+00	0,00E+00	3,18E+03	0,00E+00	3,18E+03
[Abdomen] Abdominal embolisation	0,00E+00	0,00E+00	0,00E+00	2,32E+04	0,00E+00	2,32E+04
[Abdomen] Abdominal thrombolysis	0,00E+00	0,00E+00	0,00E+00	2,27E+02	0,00E+00	2,27E+02
[Abdomen] Abdominal region biopsy	0,00E+00	0,00E+00	0,00E+00	2,55E+02	0,00E+00	2,55E+02
[Pelvis] Pelvic vessel dilatation	0,00E+00	0,00E+00	0,00E+00	1,75E+04	0,00E+00	1,75E+04
[Pelvis] Pelvic vessel embolisation	0,00E+00	0,00E+00	0,00E+00	1,67E+04	0,00E+00	1,67E+04
[Pelvis] Pelvic vessel thrombolysis	0,00E+00	0,00E+00	0,00E+00	1,76E+02	0,00E+00	1,76E+02
[Limbs] Upper limb dilatation	0,00E+00	0,00E+00	0,00E+00	1,26E+02	0,00E+00	1,26E+02
[Limbs] Upper limb embolisation	0,00E+00	0,00E+00	0,00E+00	4,81E+01	0,00E+00	4,81E+01
[Limbs] Upper limb thrombolysis	0,00E+00	0,00E+00	0,00E+00	2,20E+00	0,00E+00	2,20E+00
[Limbs] Popliteal dilatation (behind knee)	0,00E+00	0,00E+00	0,00E+00	1,12E+03	0,00E+00	1,12E+03
[Limbs] Lower limb dilatation	0,00E+00	0,00E+00	0,00E+00	5,79E+03	0,00E+00	5,79E+03
[Limbs] Lower limb embolisation	0,00E+00	0,00E+00	0,00E+00	3,19E+01	0,00E+00	3,19E+01
[Limbs] Lower limb thrombolysis	0,00E+00	0,00E+00	0,00E+00	3,66E+02	0,00E+00	3,66E+02
[Limbs] Limbs biopsy	0,00E+00	0,00E+00	0,00E+00	0,00E+00	9,00E-01	9,00E-01
[Other] Thoracic puncture	0,00E+00	0,00E+00	0,00E+00	1,49E+03	0,00E+00	1,49E+03
[Other] Abdominal puncture	0,00E+00	0,00E+00	0,00E+00	3,05E+03	0,00E+00	3,05E+03
[Other] Extremities puncture	0,00E+00	0,00E+00	0,00E+00	1,31E+02	0,00E+00	1,31E+02
[Other] Vertebroplasty	0,00E+00	0,00E+00	0,00E+00	1,59E+03	0,00E+00	1,59E+03
Total	0,00E+00	1,35E+01	0,00E+00	5,28E+05	9,00E-01	5,28E+05

Table 4: Annual collective dose (in mSv) delivered through diagnostic interventions by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Blood vessels] Cerebral angiography	0,00E+00	0,00E+00	0,00E+00	1,66E+04	0,00E+00	1,66E+04
[Blood vessels] Petrous phlebography	0,00E+00	0,00E+00	0,00E+00	4,82E+01	0,00E+00	4,82E+01
[Blood vessels] Coronary angiography	0,00E+00	0,00E+00	0,00E+00	3,83E+05	4,48E+01	3,83E+05
[Blood vessels] Thoracic aortography	0,00E+00	0,00E+00	0,00E+00	7,16E+03	0,00E+00	7,16E+03
[Blood vessels] Bronchial arteriography	0,00E+00	0,00E+00	0,00E+00	3,52E+03	0,00E+00	3,52E+03
[Blood vessels] Pulmonary arteriography	0,00E+00	0,00E+00	0,00E+00	5,56E+02	0,00E+00	5,56E+02
[Blood vessels] Upper venacavography	0,00E+00	0,00E+00	0,00E+00	5,55E+03	0,00E+00	5,55E+03
[Blood vessels] Abdominal aortography	0,00E+00	0,00E+00	0,00E+00	2,73E+04	0,00E+00	2,73E+04
[Blood vessels] Renal arteriography	0,00E+00	0,00E+00	0,00E+00	9,17E+03	0,00E+00	9,17E+03
[Blood vessels] Mesenteric arteriography	0,00E+00	0,00E+00	0,00E+00	3,19E+03	0,00E+00	3,19E+03
[Blood vessels] Lower venacavography	0,00E+00	0,00E+00	0,00E+00	3,82E+03	0,00E+00	3,82E+03
[Blood vessels] Renal phlebography	0,00E+00	0,00E+00	0,00E+00	1,65E+02	0,00E+00	1,65E+02
[Blood vessels] Suprarenal phlebography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Blood vessels] Pelvic arteriography	0,00E+00	0,00E+00	0,00E+00	2,34E+04	0,00E+00	2,34E+04
[Blood vessels] Ovarian phlebography	0,00E+00	0,00E+00	0,00E+00	7,06E+01	0,00E+00	7,06E+01
[Blood vessels] Spermatic phlebography	0,00E+00	0,00E+00	0,00E+00	1,56E+02	0,00E+00	1,56E+02
[Blood vessels] Upper limb arteriography	0,00E+00	0,00E+00	0,00E+00	7,64E+03	0,00E+00	7,64E+03
[Blood vessels] Upper limb phlebography	0,00E+00	0,00E+00	0,00E+00	5,79E+03	1,60E+01	5,81E+03
[Blood vessels] Lower limb arteriography	0,00E+00	0,00E+00	0,00E+00	5,10E+04	0,00E+00	5,10E+04
[Blood vessels] Lower limb phlebography	0,00E+00	0,00E+00	0,00E+00	5,06E+03	1,92E+02	5,26E+03
[Lymph vessels] Thoracic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Abdominal lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00

[Lymph vessels] Pelvic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Upper limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Lower limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Total	0,00E+00	0,00E+00	0,00E+00	5,53E+05	2,53E+02	5,53E+05

Table 5: Annual collective dose (in mSv) delivered through CT examinations by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Head] Orbits	0,00E+00	0,00E+00	0,00E+00	1,77E+04	1,46E+03	1,92E+04
[Head] Temporal bones	0,00E+00	0,00E+00	0,00E+00	2,19E+04	5,66E+02	2,25E+04
[Head] Temporo-mandibular joint	0,00E+00	0,00E+00	0,00E+00	2,90E+03	3,92E+02	3,30E+03
[Head] Sella turcica	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Face	0,00E+00	0,00E+00	0,00E+00	3,76E+03	1,09E+03	4,85E+03
[Head] Dental	5,74E+03	0,00E+00	0,00E+00	5,51E+02	1,36E+03	7,65E+03
[Head] Cerebrum	0,00E+00	0,00E+00	0,00E+00	1,61E+05	1,68E+04	1,78E+05
[Head] Posterior fossa	0,00E+00	0,00E+00	0,00E+00	5,87E+03	1,23E+02	5,99E+03
[Head] Brain vascular	0,00E+00	0,00E+00	0,00E+00	4,61E+03	1,08E+03	5,69E+03
[Head] Pituitary gland	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Sinuses	0,00E+00	0,00E+00	0,00E+00	4,10E+03	3,51E+03	7,61E+03
[Head] Internal auditory meatus	0,00E+00	0,00E+00	0,00E+00	7,33E+02	5,29E+00	7,39E+02
[Head] Nasal cavity	0,00E+00	0,00E+00	0,00E+00	2,76E+03	4,49E+02	3,21E+03
[Head] Mouth	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Neck] Cervical spine	0,00E+00	0,00E+00	0,00E+00	4,98E+04	6,14E+03	5,59E+04
[Neck] Neck	0,00E+00	0,00E+00	0,00E+00	4,25E+04	8,03E+03	5,06E+04
[Neck] Larynx	0,00E+00	0,00E+00	0,00E+00	8,46E+02	5,44E+02	1,39E+03
[Neck] Pharynx	0,00E+00	0,00E+00	0,00E+00	1,07E+02	0,00E+00	1,07E+02
[Neck] Neck vascular	0,00E+00	0,00E+00	0,00E+00	1,70E+04	7,74E+02	1,77E+04
[Chest + abdomen] Thoracic spine	0,00E+00	0,00E+00	0,00E+00	3,66E+04	3,57E+03	4,02E+04
[Chest + abdomen] Mediastinum	0,00E+00	0,00E+00	0,00E+00	4,51E+04	9,51E+01	4,52E+04
[Chest + abdomen] Lungs standard	0,00E+00	0,00E+00	0,00E+00	4,56E+05	8,98E+04	5,46E+05

[Chest + abdomen] Lungs High Resolution	0,00E+00	0,00E+00	0,00E+00	1,15E+04	9,68E+03	2,12E+04
[Chest + abdomen] Heart	0,00E+00	0,00E+00	0,00E+00	3,90E+04	6,51E+03	4,55E+04
[Chest + abdomen] Thoracic aorta	0,00E+00	0,00E+00	0,00E+00	2,91E+04	3,79E+03	3,29E+04
[Chest + abdomen] Lungs vascular	0,00E+00	0,00E+00	0,00E+00	4,57E+04	5,48E+03	5,12E+04
[Abdomen] Lumbar spine	0,00E+00	0,00E+00	0,00E+00	2,12E+05	6,75E+04	2,80E+05
[Abdomen] Full abdomen	0,00E+00	0,00E+00	0,00E+00	1,55E+06	2,26E+05	1,77E+06
[Abdomen] Upper abdomen	0,00E+00	0,00E+00	0,00E+00	1,35E+05	1,09E+05	2,44E+05
[Abdomen] Liver / pancreas	0,00E+00	0,00E+00	0,00E+00	3,41E+04	1,22E+03	3,54E+04
[Abdomen] Kidneys / Supra-renal glands	0,00E+00	0,00E+00	0,00E+00	3,66E+04	2,28E+03	3,88E+04
[Abdomen] Enteroclysis	0,00E+00	0,00E+00	0,00E+00	3,56E+02	1,36E+02	4,92E+02
[Pelvis] Hip / pelvic bone	0,00E+00	0,00E+00	0,00E+00	3,87E+04	2,84E+04	6,73E+04
[Pelvis] Sacrum / coccyx	0,00E+00	0,00E+00	0,00E+00	1,68E+04	1,53E+03	1,83E+04
[Pelvis] Sacro-iliac joint	0,00E+00	0,00E+00	0,00E+00	9,27E+03	2,36E+04	3,29E+04
[Pelvis] Pelvimetry (obstetric)	0,00E+00	0,00E+00	0,00E+00	1,67E+03	1,63E+01	1,68E+03
[Pelvis] Pelvis (soft tissues/vascular)	0,00E+00	0,00E+00	0,00E+00	5,44E+04	1,05E+04	6,50E+04
[Pelvis] Virtual CT colonoscopy	0,00E+00	0,00E+00	0,00E+00	3,79E+03	5,77E+03	9,56E+03
[Neck + chest + abdomen] Full spine	0,00E+00	0,00E+00	0,00E+00	3,79E+05	9,14E+03	3,88E+05
[Chest + abdomen] Chest & abdomen	0,00E+00	0,00E+00	0,00E+00	2,78E+05	1,10E+04	2,89E+05
[Abdomen + pelvis] Abdomen & pelvis	0,00E+00	0,00E+00	0,00E+00	4,31E+05	9,30E+03	4,40E+05
[Abdomen + pelvis] CT urography	0,00E+00	0,00E+00	0,00E+00	1,38E+04	2,48E+03	1,63E+04
[Chest + abdomen + pelvis] Whole trunk	0,00E+00	0,00E+00	0,00E+00	6,45E+04	0,00E+00	6,45E+04
[Limbs] Shoulder	0,00E+00	0,00E+00	0,00E+00	1,06E+04	3,35E+03	1,40E+04
[Limbs] Elbow	0,00E+00	0,00E+00	0,00E+00	3,86E+03	7,24E+02	4,62E+03
[Limbs] Wrist	0,00E+00	0,00E+00	0,00E+00	9,54E+03	1,98E+03	1,16E+04
[Limbs] Hand	0,00E+00	0,00E+00	0,00E+00	2,69E+03	4,00E+02	3,15E+03

[Limbs] Thigh	0,00E+00	0,00E+00	0,00E+00	3,54E+02	2,82E+01	3,82E+02
[Limbs] Knee	0,00E+00	0,00E+00	0,00E+00	7,01E+02	2,09E+02	9,10E+02
[Limbs] Lower leg	0,00E+00	0,00E+00	0,00E+00	1,39E+02	5,02E+01	1,89E+02
[Limbs] Calcaneum	0,00E+00	0,00E+00	0,00E+00	6,78E+01	2,00E+01	8,84E+01
[Limbs] Ankle	0,00E+00	0,00E+00	0,00E+00	5,37E+02	8,95E+01	6,34E+02
[Limbs] Foot	0,00E+00	0,00E+00	0,00E+00	3,33E+02	1,16E+02	4,55E+02
[Other] CT angiography	0,00E+00	0,00E+00	0,00E+00	4,70E+05	2,79E+05	7,49E+05
[Other] CT intervention	0,00E+00	0,00E+00	0,00E+00	3,79E+05	5,57E+04	4,35E+05
Total	5,74E+03	0,00E+00	0,00E+00	5,13E+06	1,01E+06	6,15E+06

Table 6: Annual collective dose (in mSv) delivered through dental examinations by different healthcare providers

Dental examinations	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Teeth & gums] Total Periapical	1,24E+04	9,00E+00	0,00E+00	1,73E+02	0,00E+00	1,25E+04
[Teeth & gums] Total Bitewing	2,29E+04	3,58E+00	0,00E+00	1,90E+02	0,00E+00	2,31E+04
[Teeth & gums] Occlusal	1,01E+03	2,04E+00	0,00E+00	5,49E+00	6,18E-01	1,02E+03
[Teeth & gums] Panoramic radiography (OPG)	8,39E+03	1,82E+02	0,00E+00	1,31E+03	1,32E+02	1,00E+04
[Teeth & gums] CT (mandible/maxilla)	6,08E+03	1,04E+03	0,00E+00	4,65E+02	3,33E+02	7,93E+03
[Head] Skull	1,95E+03	2,16E+01	0,00E+00	2,99E+02	6,74E+00	2,28E+03
[Head] Teleradiography	4,94E+03	5,46E+01	0,00E+00	7,28E+02	4,35E+01	5,76E+03
[Limbs] Hand	1,48E+00	1,44E-03	0,00E+00	1,07E+00	2,04E-01	2,75E+00
Total	5,76E+04	1,32E+03	0,00E+00	3,17E+03	5,16E+02	6,26E+04

Table 7: Annual collective dose (in mSv) delivered through mammography examinations by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Breast] Symptomatic (1-2 views / 1 or both breasts)	0,00E+00	3,86E+03	0,00E+00	2,65E+04	1,51E+04	4,55E+04
[Breast] Screening (1-2 views / both breasts)	0,00E+00	9,11E+02	0,00E+00	1,16E+04	3,90E+03	1,64E+04
Total	0,00E+00	4,77E+03	0,00E+00	3,81E+04	1,90E+04	6,19E+04

Table 8: Annual collective dose (in mSv) delivered through bone densitometry examinations by different healthcare providers

	Dental Practices	Medical Practices	Chiropractors	Hospital departments	Radiology institutes	Total
[Abdomen] Lumbar spine	0,00E+00	4,19E+00	0,00E+00	1,36E+02	3,56E+01	1,75E+02
[Pelvis] Hip	0,00E+00	3,12E+00	0,00E+00	1,01E+02	2,66E+01	1,31E+02
[Limbs] Upper or lower limb	0,00E+00	6,33E-02	0,00E+00	2,05E+00	7,63E-01	2,87E+00
[Other] Whole body	0,00E+00	0,00E+00	0,00E+00	3,79E+00	0,00E+00	3,79E+00
Total	0,00E+00	7,37E+00	0,00E+00	2,42E+02	6,30E+01	3,13E+02

Appendix 9:

Number of examinations (N),
Collective dose (S), Frequency of
examinations (N/1000
population), and average effective
dose (E/caput)

Table 1: Frequency and dosimetry results (radiography examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Head] Orbits	2,75E+04	8,25E+02	3,57E+00	1,07E-04
[Head] Temporal bones	8,78E+03	5,27E+02	1,14E+00	6,84E-05
[Head] Sphenoid bone	1,38E+04	4,13E+02	1,79E+00	5,36E-05
[Head] Nose	1,03E+04	4,13E+02	1,34E+00	5,36E-05
[Head] Sinuses	7,53E+04	3,47E+03	9,78E+00	4,50E-04
[Head] Zygomas	6,88E+02	2,75E+01	8,93E-02	3,57E-06
[Head] Temporo-mandibular joint	1,63E+03	6,51E+01	2,12E-01	8,45E-06
[Head] Cervico-occipital hinge	2,46E+03	1,03E+02	3,19E-01	1,34E-05
[Head] Maxilla	1,29E+03	5,17E+01	1,67E-01	6,72E-06
[Head] Mandible	7,53E+03	3,01E+02	9,78E-01	3,91E-05
[Head] Cephalometry	1,97E+02	7,88E+00	2,56E-02	1,02E-06
[Head] Dacryocystography	2,94E+03	2,36E+02	3,82E-01	3,06E-05
[Head] Sialography	1,11E+02	4,43E+00	1,44E-02	5,75E-07
[Head] Eyes	6,50E+02	2,60E+01	8,44E-02	3,37E-06
[Neck] Cervical spine	1,95E+05	1,99E+04	2,53E+01	2,58E-03
[Neck] Larynx	6,89E+03	3,45E+02	8,95E-01	4,47E-05
[Neck] Pharynx	6,99E+03	3,49E+02	9,08E-01	4,54E-05
[Neck] Trachea	6,25E+03	6,25E+02	8,11E-01	8,11E-05
[Chest / Thorax] Thoracic spine	8,62E+04	3,75E+04	1,12E+01	4,87E-03
[Chest / Thorax] Shoulder blades / scapulae	2,07E+05	6,00E+03	2,69E+01	7,79E-04
[Chest / Thorax] Collar bone(s) / clavicle(s)	4,05E+04	1,17E+03	5,26E+00	1,52E-04
[Chest / Thorax] Acromio-clavicular joint	3,94E+04	1,14E+03	5,12E+00	1,48E-04
[Chest / Thorax] Sterno-clavicular joint	6,00E+03	1,74E+02	7,79E-01	2,26E-05

[Chest / Thorax] Manubrio-sternal joint	7,67E+01	2,22E+00	9,96E-03	2,89E-07
[Chest / Thorax] Sternum	3,45E+03	1,00E+02	4,48E-01	1,30E-05
[Chest / Thorax] Ribs	3,55E+04	1,78E+03	4,61E+00	2,31E-04
[Chest / Thorax] Lung	1,75E+06	8,84E+04	2,27E+02	1,15E-02
[Chest / Thorax] Thoracic inlet	7,34E+03	2,79E+02	9,53E-01	3,62E-05
[Chest / Thorax] Bronchography	5,34E+02	2,03E+01	6,93E-02	2,63E-06
[Abdomen] Lumbar spine	3,40E+05	5,76E+05	4,41E+01	7,48E-02
[Abdomen] Lumbo-sacral joint	4,16E+03	3,42E+03	5,40E-01	4,44E-04
[Abdomen] Abdomen (plain film, patient supine or erect)	1,51E+05	1,18E+05	1,96E+01	1,53E-02
[Pelvis] Ilium / ischium / pubis	2,56E+05	2,08E+05	3,32E+01	2,70E-02
[Pelvis] Sacrum	1,49E+04	1,61E+04	1,93E+00	2,09E-03
[Pelvis] Sacro-iliac joint	1,41E+04	1,53E+04	1,83E+00	1,98E-03
[Pelvis] Coccyx	5,95E+03	6,42E+03	7,73E-01	8,34E-04
[Pelvis] Pelvimetry (obstetric)	4,45E+01	2,40E+01	5,78E-03	3,12E-06
[Pelvis] Hips (one or both)	1,81E+05	2,05E+05	2,35E+01	2,66E-02
[Pelvis] Pelvis (soft tissue)	8,16E+03	4,41E+03	1,06E+00	5,72E-04
[Limbs] Upper arm (humerus)	4,68E+04	2,86E+02	6,08E+00	3,71E-05
[Limbs] Elbow	1,10E+05	6,62E+01	1,43E+01	8,59E-06
[Limbs] Forearm (radius & ulna)	5,66E+04	6,22E+01	7,35E+00	8,08E-06
[Limbs] Wrist (scaphoid)	2,74E+05	5,49E+01	3,56E+01	7,13E-06
[Limbs] Hand	2,25E+05	4,50E+01	2,92E+01	5,84E-06
[Limbs] Fingers & thumbs	2,29E+05	2,29E+01	2,97E+01	2,97E-06
[Limbs] Femur	3,45E+04	1,38E+03	4,48E+00	1,79E-04
[Limbs] Knee joint	5,45E+05	4,36E+02	7,08E+01	5,66E-05
[Limbs] Knee cap (patella)	5,55E+04	2,22E+01	7,21E+00	2,88E-06

[Limbs] Lower leg (tibia & fibula)	6,41E+04	2,56E+01	8,32E+00	3,33E-06
[Limbs] Ankle	3,41E+05	1,36E+02	4,43E+01	1,77E-05
[Limbs] Foot	3,38E+05	1,35E+02	4,39E+01	1,76E-05
[Limbs] Calcaneum (heel)	4,06E+04	1,62E+01	5,27E+00	2,11E-06
[Limbs] Toes	8,68E+04	3,47E+01	1,13E+01	4,51E-06
[Limbs] Whole leg (Leg length)	1,59E+04	3,19E+02	2,06E+00	4,14E-05
[Trunk] Scoliosis (Whole spine)	1,40E+04	4,75E+03	1,82E+00	6,17E-04
[Head & trunk] Whole skeleton (Skeletal survey)	1,96E+03	6,75E+03	2,54E-01	8,77E-04
Total	6,00E+06	1,33E+06	7,79E+02	1,73E-01

Table 2: Frequency and dosimetry results (fluoroscopy examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[GI tract] Oesophagus	1,45E+04	1,59E+04	1,88E+00	2,07E-03
[GI tract] Stomach & duodenum	9,07E+03	1,09E+05	1,18E+00	1,41E-02
[GI tract] Small intestine	2,50E+03	9,52E+03	3,25E-01	1,24E-03
[GI tract] Enteroclysis	7,07E+02	4,95E+03	9,18E-02	6,42E-04
[GI tract] Colon	5,25E+03	6,30E+04	6,82E-01	8,18E-03
[GI tract] Defecography	2,04E+03	2,45E+04	2,65E-01	3,18E-03
[Biliary tract] Retrograde cholangiography	1,72E+03	3,10E+03	2,23E-01	4,02E-04
[Biliary tract] Operative cholangiography	8,82E+02	1,59E+03	1,15E-01	2,06E-04
[Biliary tract] Intravenous cholangiography	2,72E+02	4,90E+02	3,53E-02	6,36E-05
[Biliary tract] T drain cholangiography	9,60E+02	2,50E+03	1,25E-01	3,24E-04
[Biliary tract] Transhepatic cholangiography	3,23E+02	2,61E+03	4,19E-02	3,39E-04
[Biliary tract] ERCP	5,33E+03	2,93E+04	6,92E-01	3,80E-03
[Biliary tract] Retrograde pancreatography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Biliary tract] Cholecystography	2,52E+00	7,81E+00	3,27E-04	1,01E-06
[Uro-genital] Intravenous urography	8,81E+03	1,85E+04	1,14E+00	2,40E-03
[Uro-genital] Retrograde pyelography	7,17E+03	1,65E+04	9,31E-01	2,14E-03
[Uro-genital] Nephrostography	1,88E+03	3,01E+03	2,44E-01	3,90E-04
[Uro-genital] Retrograde cystography	6,15E+03	1,11E+04	7,99E-01	1,44E-03
[Uro-genital] Micturitional cysto-urethrography	9,83E+03	3,44E+04	1,28E+00	4,47E-03
[Uro-genital] Urethrography	2,31E+03	2,54E+03	3,00E-01	3,30E-04
[Uro-genital] Hysterosalpingography	4,74E+03	1,19E+04	6,15E-01	1,54E-03
[Spinal cord] Cervical myelography	2,24E+02	1,34E+02	2,91E-02	1,74E-05
[Spinal cord] Thoracic myelography	2,77E+02	1,08E+03	3,60E-02	1,40E-04

[Spinal cord] Lumbar myelography	3,41E+03	2,21E+04	4,43E-01	2,88E-03
[Spinal cord] Sacral myelography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Spinal cord] Whole spine myelography	5,25E+01	5,77E+02	6,82E-03	7,49E-05
[Joints] Temporal-mandibular joint arthrography	7,75E+01	6,03E+01	1,01E-02	7,83E-06
[Joints] Shoulder arthrography	4,08E+04	5,31E+03	5,30E+00	6,89E-04
[Joints] Hip arthrography	6,36E+03	2,03E+02	8,26E-01	2,64E-05
[Joints] Elbow arthrography	4,84E+02	2,91E+00	6,28E-02	3,77E-07
[Joints] Wrist arthrography	2,66E+03	2,66E+00	3,45E-01	3,46E-07
[Joints] Knee arthrography	4,21E+03	3,67E+02	5,47E-01	4,76E-05
[Joints] Ankle arthrography	1,22E+03	3,67E+00	1,58E-01	4,76E-07
[Other] Colonoscopy	1,18E+03	5,71E+03	1,53E-01	7,41E-04
[Other] Chest fluoroscopy	2,71E+03	8,65E+03	3,52E-01	1,12E-03
[Other] Abdominal fluoroscopy	1,44E+03	5,80E+03	1,87E-01	7,53E-04
[Other] Limbs/Joints fluoroscopy	3,09E+03	6,42E+02	4,01E-01	8,34E-05
Total	1,53E+05	4,15E+05	1,99E+01	5,39E-02

Table 3: Frequency and dosimetry results (therapeutic interventions)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Head & neck] Cerebral dilatation/stenting	3,10E+02	5,11E+03	4,02E-02	6,63E-04
[Head & neck] Cerebral embolisation (AVM, a.aneurysm, a.tumour)	4,38E+02	1,44E+04	5,68E-02	1,87E-03
[Head & neck] Cerebral thrombolysis	2,75E+02	4,65E+02	3,57E-02	6,04E-05
[Head & neck] Head & neck puncture	4,48E+02	7,61E+01	5,81E-02	9,88E-06
[Chest] Coronary dilatation/stenting (PTCA)	1,82E+04	3,10E+05	2,37E+00	4,03E-02
[Chest] Cardiac pacemaker fitting (temporary or permanent)	2,60E+03	7,29E+03	3,38E-01	9,46E-04
[Chest] Central venous line fitting (Hickman line)	3,93E+02	9,83E+01	5,11E-02	1,28E-05
[Chest] Cardiac thermo-ablation	7,72E+02	1,57E+04	1,00E-01	2,04E-03
[Chest] Valvuloplasty	1,37E+02	4,03E+03	1,78E-02	5,23E-04
[Chest] Inferior vena cava (IVC) filter fitting	2,64E+02	9,50E+02	3,43E-02	1,23E-04
[Chest] Oesophagus dilatation/stenting	3,43E+02	5,15E+02	4,46E-02	6,69E-05
[Chest] Thoracic dilatation/stenting	3,89E+02	1,14E+04	5,05E-02	1,49E-03
[Chest] Thoracic embolisation	1,56E+01	4,58E+02	2,02E-03	5,95E-05
[Chest] Thoracic thrombolysis	2,21E+01	6,50E+02	2,87E-03	8,44E-05
[Chest] Thoracic region biopsy	6,16E+01	2,46E+01	8,00E-03	3,20E-06
[Chest] Electrophysiology	1,10E+03	3,64E+03	1,43E-01	4,73E-04
[Abdomen] Bile duct dilatation/stenting	6,01E+02	8,41E+03	7,80E-02	1,09E-03
[Abdomen] Bile duct drainage	1,06E+03	4,04E+04	1,37E-01	5,25E-03
[Abdomen] Bile duct stone extraction	2,05E+02	1,43E+03	2,66E-02	1,86E-04
[Abdomen] Renal artery dilatation/stenting	3,17E+02	3,81E+03	4,12E-02	4,94E-04
[Abdomen] Renal drainage	5,48E+02	1,03E+04	7,11E-02	1,34E-03
[Abdomen] Lithotripsy	2,95E+02	1,12E+03	3,83E-02	1,45E-04
[Abdomen] Nephrostomy	1,09E+02	1,48E+03	1,42E-02	1,92E-04

[Abdomen] TIPS (liver)	1,68E+02	1,08E+04	2,18E-02	1,41E-03
[Abdomen] Abdominal dilatation/stenting	2,12E+03	3,18E+03	2,75E-01	4,13E-04
[Abdomen] Abdominal embolisation	6,31E+02	2,32E+04	8,20E-02	3,01E-03
[Abdomen] Abdominal thrombolysis	1,37E+01	2,27E+02	1,78E-03	2,94E-05
[Abdomen] Abdominal region biopsy	1,70E+02	2,55E+02	2,20E-02	3,31E-05
[Pelvis] Pelvic vessel dilatation	1,21E+03	1,75E+04	1,57E-01	2,27E-03
[Pelvis] Pelvic vessel embolisation	4,49E+02	1,67E+04	5,83E-02	2,17E-03
[Pelvis] Pelvic vessel thrombolysis	1,55E+01	1,76E+02	2,01E-03	2,28E-05
[Limbs] Upper limb dilatation	6,98E+02	1,26E+02	9,07E-02	1,63E-05
[Limbs] Upper limb embolisation	1,20E+02	4,81E+01	1,56E-02	6,25E-06
[Limbs] Upper limb thrombolysis	1,69E+01	2,20E+00	2,20E-03	2,86E-07
[Limbs] Popliteal dilatation (behind knee)	4,80E+02	1,12E+03	6,23E-02	1,45E-04
[Limbs] Lower limb dilatation	6,44E+03	5,79E+03	8,36E-01	7,52E-04
[Limbs] Lower limb embolisation	3,55E+01	3,19E+01	4,61E-03	4,15E-06
[Limbs] Lower limb thrombolysis	4,07E+02	3,66E+02	5,28E-02	4,75E-05
[Limbs] Limbs biopsy	1,00E+00	9,00E-01	1,30E-04	1,17E-07
[Other] Thoracic puncture	6,54E+02	1,49E+03	8,49E-02	1,94E-04
[Other] Abdominal puncture	1,20E+03	3,05E+03	1,56E-01	3,96E-04
[Other] Extremities puncture	8,94E+02	1,31E+02	1,16E-01	1,71E-05
[Other] Vertebroplasty	1,26E+02	1,59E+03	1,64E-02	2,07E-04
Total	4,60E+04	5,28E+05	5,97E+00	6,85E-02

Table 4: Frequency and dosimetry results (diagnostic interventions)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Blood vessels] Cerebral angiography	3,32E+03	1,66E+04	4,31E-01	2,16E-03
[Blood vessels] Petrous phlebography	3,76E+01	4,82E+01	4,89E-03	6,25E-06
[Blood vessels] Coronary angiography	3,42E+04	3,83E+05	4,44E+00	4,97E-02
[Blood vessels] Thoracic aortography	1,43E+03	7,16E+03	1,86E-01	9,30E-04
[Blood vessels] Bronchial arteriography	7,03E+02	3,52E+03	9,13E-02	4,57E-04
[Blood vessels] Pulmonary arteriography	1,11E+02	5,56E+02	1,44E-02	7,22E-05
[Blood vessels] Upper venacavography	1,11E+03	5,55E+03	1,44E-01	7,20E-04
[Blood vessels] Abdominal aortography	2,28E+03	2,73E+04	2,96E-01	3,55E-03
[Blood vessels] Renal arteriography	1,53E+03	9,17E+03	1,98E-01	1,19E-03
[Blood vessels] Mesenteric arteriography	3,19E+02	3,19E+03	4,14E-02	4,14E-04
[Blood vessels] Lower venacavography	3,19E+02	3,82E+03	4,14E-02	4,96E-04
[Blood vessels] Renal phlebography	2,75E+01	1,65E+02	3,57E-03	2,14E-05
[Blood vessels] Suprarenal phlebography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Blood vessels] Pelvic arteriography	1,44E+03	2,34E+04	1,86E-01	3,04E-03
[Blood vessels] Ovarian phlebography	2,32E+01	7,06E+01	3,02E-03	9,17E-06
[Blood vessels] Spermatic phlebography	2,50E+01	1,56E+02	3,25E-03	2,03E-05
[Blood vessels] Upper limb arteriography	9,56E+02	7,64E+03	1,24E-01	9,92E-04
[Blood vessels] Upper limb phlebography	7,26E+02	5,81E+03	9,43E-02	7,54E-04
[Blood vessels] Lower limb arteriography	6,38E+03	5,10E+04	8,28E-01	6,62E-03
[Blood vessels] Lower limb phlebography	6,57E+02	5,26E+03	8,53E-02	6,82E-04
[Lymph vessels] Thoracic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Abdominal lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Pelvic lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Upper limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Lymph vessels] Lower limb lymphangiography	0,00E+00	0,00E+00	0,00E+00	0,00E+00
Total	5,56E+04	5,53E+05	7,21E+00	7,18E-02

Table 5: Frequency and dosimetry results (CT examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Head] Orbits	1,55E+04	1,92E+04	2,01E+00	2,49E-03
[Head] Temporal bones	2,81E+04	2,25E+04	3,65E+00	2,92E-03
[Head] Temporo-mandibular joint	4,12E+03	3,30E+03	5,35E-01	4,28E-04
[Head] Sella turcica	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Face	9,69E+03	4,85E+03	1,26E+00	6,29E-04
[Head] Dental	1,53E+04	7,65E+03	1,99E+00	9,93E-04
[Head] Cerebrum	8,29E+04	1,78E+05	1,08E+01	2,31E-02
[Head] Posterior fossa	2,80E+03	5,99E+03	3,63E-01	7,77E-04
[Head] Brain vascular	2,66E+03	5,69E+03	3,45E-01	7,39E-04
[Head] Pituitary gland	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Head] Sinuses	1,52E+04	7,61E+03	1,98E+00	9,88E-04
[Head] Internal auditory meatus	1,48E+03	7,39E+02	1,92E-01	9,59E-05
[Head] Nasal cavity	6,41E+03	3,21E+03	8,32E-01	4,16E-04
[Head] Mouth	0,00E+00	0,00E+00	0,00E+00	0,00E+00
[Neck] Cervical spine	1,94E+04	5,59E+04	2,51E+00	7,26E-03
[Neck] Neck	1,82E+04	5,06E+04	2,36E+00	6,56E-03
[Neck] Larynx	5,00E+02	1,39E+03	6,49E-02	1,80E-04
[Neck] Pharynx	3,84E+01	1,07E+02	4,99E-03	1,39E-05
[Neck] Neck vascular	5,73E+03	1,77E+04	7,44E-01	2,30E-03
[Chest + abdomen] Thoracic spine	7,44E+03	4,02E+04	9,66E-01	5,22E-03
[Chest + abdomen] Mediastinum	8,37E+03	4,52E+04	1,09E+00	5,87E-03
[Chest + abdomen] Lungs standard	1,01E+05	5,46E+05	1,31E+01	7,08E-02
[Chest + abdomen] Lungs High Resolution	3,93E+03	2,12E+04	5,10E-01	2,75E-03
[Chest + abdomen] Heart	4,89E+03	4,55E+04	6,35E-01	5,90E-03
[Chest + abdomen] Thoracic aorta	5,81E+03	3,29E+04	7,55E-01	4,27E-03
[Chest + abdomen] Lungs vascular	9,18E+03	5,12E+04	1,19E+00	6,65E-03
[Abdomen] Lumbar spine	3,04E+04	2,80E+05	3,95E+00	3,63E-02
[Abdomen] Full abdomen	1,52E+05	1,77E+06	1,97E+01	2,30E-01

[Abdomen] Upper abdomen	2,59E+04	2,44E+05	3,36E+00	3,17E-02
[Abdomen] Liver / pancreas	3,55E+03	3,54E+04	4,61E-01	4,59E-03
[Abdomen] Kidneys / Supra-renal glands	3,90E+03	3,88E+04	5,06E-01	5,04E-03
[Abdomen] Enteroclysis	4,94E+01	4,92E+02	6,41E-03	6,38E-05
[Pelvis] Hip / pelvic bone	9,35E+03	6,73E+04	1,21E+00	8,74E-03
[Pelvis] Sacrum / coccyx	2,54E+03	1,83E+04	3,30E-01	2,37E-03
[Pelvis] Sacro-iliac joint	4,57E+03	3,29E+04	5,93E-01	4,27E-03
[Pelvis] Pelvimetry (obstetric)	2,34E+02	1,68E+03	3,04E-02	2,19E-04
[Pelvis] Pelvis (soft tissues/vascular)	6,60E+03	6,50E+04	8,57E-01	8,44E-03
[Pelvis] Virtual CT colonoscopy	1,08E+03	9,56E+03	1,40E-01	1,24E-03
[Neck + chest + abdomen] Full spine	2,32E+04	3,88E+05	3,02E+00	5,04E-02
[Chest + abdomen] Chest & abdomen	2,68E+04	2,89E+05	3,48E+00	3,76E-02
[Abdomen + pelvis] Abdomen & pelvis	3,76E+04	4,40E+05	4,88E+00	5,72E-02
[Abdomen + pelvis] CT urography	2,54E+03	1,63E+04	3,30E-01	2,11E-03
[Chest + abdomen + pelvis] Whole trunk	5,66E+03	6,45E+04	7,35E-01	8,38E-03
[Limbs] Shoulder	7,79E+03	1,40E+04	1,01E+00	1,82E-03
[Limbs] Elbow	2,57E+03	4,62E+03	3,33E-01	5,99E-04
[Limbs] Wrist	6,42E+03	1,16E+04	8,33E-01	1,50E-03
[Limbs] Hand	1,75E+03	3,15E+03	2,27E-01	4,09E-04
[Limbs] Thigh	3,82E+03	3,82E+02	4,97E-01	4,97E-05
[Limbs] Knee	9,10E+03	9,10E+02	1,18E+00	1,18E-04
[Limbs] Lower leg	1,89E+03	1,89E+02	2,46E-01	2,46E-05
[Limbs] Calcaneum	8,84E+02	8,84E+01	1,15E-01	1,15E-05
[Limbs] Ankle	6,34E+03	6,34E+02	8,23E-01	8,23E-05
[Limbs] Foot	4,55E+03	4,55E+02	5,91E-01	5,91E-05
[Other] CT angiography	1,92E+04	7,49E+05	2,49E+00	9,73E-02
[Other] CT intervention	1,11E+04	4,35E+05	1,45E+00	5,65E-02
Total	7,80E+05	6,15E+06	1,01E+02	7,99E-01

Table 6: Frequency and dosimetry results (dental examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Teeth & gums] Total Periapical	2,51E+06	1,25E+04	3,26E+02	1,63E-03
[Teeth & gums] Total Bitewing	2,31E+06	2,31E+04	2,99E+02	2,99E-03
[Teeth & gums] Occlusal	4,06E+04	1,02E+03	5,28E+00	1,32E-04
[Teeth & gums] Panoramic radiography (OPG)	4,17E+05	1,00E+04	5,42E+01	1,30E-03
[Teeth & gums] CT (mandible/maxilla)	1,59E+04	7,93E+03	2,06E+00	1,03E-03
[Head] Skull	3,74E+04	2,28E+03	4,86E+00	2,96E-04
[Head] Teleradiography	9,45E+04	5,76E+03	1,23E+01	7,48E-04
[Limbs] Hand	9,18E+03	2,75E+00	1,19E+00	3,57E-07
Total	5,43E+06	6,26E+04	7,05E+02	8,13E-03

Table 7: Frequency and dosimetry results (mammography examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Breast] Symptomatic (1-2 views / 1 or both breasts)	2,84E+05	4,55E+04	3,69E+01	5,90E-03
[Breast] Screening (1-2 views / both breasts)	1,02E+05	1,64E+04	1,33E+01	2,13E-03
Total	3,87E+05	6,19E+04	5,02E+01	8,03E-03

Table 8: Frequency and dosimetry results (bone densitometry examinations)

	N	S (man·mSv)	N/1000	E (mSv)/caput
[Abdomen] Lumbar spine	4,38E+04	1,75E+02	5,69E+00	2,28E-05
[Pelvis] Hip	4,35E+04	1,31E+02	5,65E+00	1,70E-05
[Limbs] Upper or lower limb	2,87E+04	2,87E+00	3,73E+00	3,73E-07
[Other] Whole body	1,13E+03	3,79E+00	1,47E-01	4,92E-07
Total	1,17E+05	3,13E+02	1,52E+01	4,06E-05

